Hard cheese



Hard cheese has a moisture content of less than 50% due to the cheese being the cheese to lose some of its moisture content and have a stronger flavour. Example: Romano

Blue cheese



Cow, sheep or goats milk with a blue or green-blue mold. The mold is derived from spores from Penicillium roqueforti, Penicillium glaucum or other being injected into the cheese curds. People who are allergic to penicillin are not advised to eat blue cheese. Example: Roquefort

Fresh, un ripened or infant cheese



Fresh cheese is not ripened, aged or fermented during the manufacturing process or at any point during the lifespan of the cheese. Fresh cheese has a very short shelf life. Example: Cottage cheese, Cream cheese.

light or lite cheese



Light cheese is made by reducing the amount of butterfat which makes the cheese rubbery in texture and much less flavourful than full fat versions of cheese. Light cheese has a high moisture content which makes it have a shorter shelf life. Example: Cheese with 7% Milk Fat, Cheddar which 19% Milk fat.

Processed cheese



This cheese is created by melting together blend of grated cheese, milk, milk solids or water, food colouring and seasonings. Example: Processed cheese shies, cheese spreads "swokies".

from milk, skimmed milk or flavoured milk. For the preparation of yoghurt, the milk should be free from contamination. The solid content (not fat should be between 11–15% which can be obtained by adding skin or whole milk powder in fresh milk that normally contains 8% solids. the product can be further improved by adding small amount of modified gums which bind water and impart thickening to the product. At this stage the size of the fat particles in the milk should be around 2µm because this improves the milk's viscosity, product's stability and milk appear form. The milk is then heated at 80–90°C for 30 min., starter culture is added to it. Heating improves the milk by inactivating immunoglobulins, remove excessive oxygen to produce micro aerophilic environment which support the growth of starter culture. Besides, heating also induce the interactions between whey or serum proteins and casein which increase yoghurt viscosity. The milk is now cooled to 40–43°C so as to allow fermentation using starter organisms such as *Streptococcus salivarius* sub *sp. thermophilus* and *Lactobacillus delbruckii* sub *sp.* bulgaricus together at a level of 2% by volume (10^6–10^7 cfu/ml). It is to be carried out for about 4h during which lactose is converted into

TN_GOVT_XII_Micro_Biology_CH05.indd 81 04-03-2019 10:49:29