3. $ye^{\frac{x}{y}}dx=\left(xe^{\frac{x}{y}}+y\right)dy$

4. $2xydx+\left(x^2+2y^2\right)dy=0$

5. $\left(y^2-2xy\right)dx=\left(x^2-2xy\right)dy$

6. $x\frac{dy}{dx}=y-x\cos^2\left(\frac{y}{x}\right)$

7. $\left(1+3e^{\frac{y}{x}}\right)dy+3e^{\frac{y}{x}}\left(1-\frac{y}{x}\right)dx=0$, given that $y=0$ when $x=1$

8. $\left(x^2+y^2\right)dy=xy\,dx$. It is given that $y(1)=1$ and $y(x_0)=e$. Find the value of x_0.

10.7 First Order Linear Differential Equations

A **first order differential equation** of the form

$$\frac{dy}{dx}+Py=Q. \qquad \ldots (1)$$

where P and Q are functions of x only. Here no product of y and its derivative $\frac{dy}{dx}$ occur and the dependent variable y and its derivative with respect to independent variable x occurs only in the first degree.

To integrate (1), let us consider the homogeneous equation $\frac{dy}{dx}+Py=0$. ...(2)

The equation (2) can be integrated as follows:



Separating the variables, $\frac{dy}{y}=-Pdx$.

On integration, we get $ye^{\int Pdx}=C$.

Now, $\frac{d}{dx}\left(ye^{\int Pdx}\right)=e^{\int Pdx}\frac{dy}{dx}+y.Pe^{\int Pdx}$

$$=e^{\int Pdx}\left(\frac{dy}{dx}+Py\right)=Qe^{\int Pdx} \qquad \ldots (3)\ (\text{using } (1))$$

Integrating both sides of (3) with respect to x, we get the solution of the given differential equation as

$$ye^{\int Pdx}=\int Qe^{\int Pdx}dx+C.$$

Here $e^{\int Pdx}$ is known as the **integrating factor (I.F.)** of (1).

Remarks

1. The solution of linear differential equation is
   $y\times(I.F)=\int Q(I.F)dx+C$, where C is an arbitrary constant.
2. In the integrating factor $e^{\int Pdx}$, P is the coefficient of y in the differential equation provided the coefficient of $\frac{dy}{dx}$ is unity.

Chapter 10 Differential Equation.indd 166 26-02-2020 17:25:27