11. The function $\sin^4 x + \cos^4 x$ is increasing in the interval

(1) $\left[\frac{5\pi}{8}, \frac{3\pi}{4}\right]$ (2) $\left[\frac{\pi}{2}, \frac{5\pi}{8}\right]$ (3) $\left[\frac{\pi}{4}, \frac{\pi}{2}\right]$ (4) $\left[0, \frac{\pi}{4}\right]$

12. The number given by the Rolle's theorem for the function $x^3 - 3x^2, x \in [0,3]$ is

(1) 1 (2) $\sqrt{2}$ (3) $\frac{3}{2}$ (4) 2

13. The number given by the Mean value theorem for the function $\frac{1}{x}$, $x \in [1,9]$ is

(1) 2 (2) 2.5 (3) 3 (4) 3.5

14. The minimum value of the function $|3-x|+9$ is

(1) 0 (2) 3 (3) 6 (4) 9

15. The maximum slope of the tangent to the curve $y = e^x \sin x, x \in [0, 2\pi]$ is at

(1) $x = \frac{\pi}{4}$ (2) $x = \frac{\pi}{2}$ (3) $x = \pi$ (4) $x = \frac{3\pi}{2}$

16. The maximum value of the function $x^2 e^{-2x}, x > 0$ is

(1) $\frac{1}{e}$ (2) $\frac{1}{2e}$ (3) $\frac{1}{e^2}$ (4) $\frac{4}{e^4}$

17. One of the closest points on the curve $x^2 - y^2 = 4$ to the point $(6,0)$ is

(1) $(2,0)$ (2) $\left(\sqrt{5},1\right)$ (3) $\left(3,\sqrt{5}\right)$ (4) $\left(\sqrt{13},-\sqrt{3}\right)$

18. The maximum value of the product of two positive numbers, when their sum of the squares is 200, is

(1) 100 (2) $25\sqrt{7}$ (3) 28 (4) $24\sqrt{14}$

19. The curve $y = ax^4 + bx^2$ with $ab > 0$

(1) has no horizontal tangent (2) is concave up

(3) is concave down (4) has no points of inflection

20. The point of inflection of the curve $y = (x-1)^3$ is

(1) $(0,0)$ (2) $(0,1)$ (3) $(1,0)$ (4) $(1,1)$

Chapter 7 Differential Calculus Original-new.indd 55 26-02-2020 17:28:06