


Figure 5.7: Meristem Culture

3. Protoplast Culture:

Protoplasts are cells without a cell wall, but bound by a cell membrane or plasma membrane. Using protoplasts, it is possible to regenerate whole plants from single cells and also develop somatic hybrids. The steps involved in protoplast culture.

i. **Isolation of protoplast:** Small bits of plant tissue like leaf tissue are used for isolation of protoplast. The leaf tissue is immersed in 0.5% Macrozyme and 2% Onozuka cellulase enzymes dissolved in 13% sorbitol or mannitol at pH 5.4. It is then incubated over-night at 25°C. After a gentle teasing of cells, protoplasts are obtained, and these are then transferred to 20% sucrose solution to retain their viability. They are then centrifuged to get pure protoplasts as different from debris of cell walls.

ii. **Fusion of protoplast:** It is done through the use of a suitable fusogen. This is normally PEG (Polyethylene Glycol). The isolated protoplast are incubated in 25 to 30% concentration of PEG with Ca++ ions and the protoplast shows agglutination (the formation of clumps of cells) and fusion.

iii. **Culture of protoplast:** MS liquid medium is used with some modification in droplet, plating or micro-drop array techniques. Protoplast viability is tested with fluorescein diacetate before the culture. The cultures are incubated in continuous light 1000-2000 lux at 25°C. The cell wall formation occurs within 24-48 hours and the first division of new cells occurs between 2-7 days of culture.

iv. **Selection of somatic hybrid cells:** The fusion product of protoplasts without nucleus of different cells is called a cybrid. Following this nuclear fusion take place. This process is called somatic hybridization.




Figure 5.8: Protoplast Culture

4. Cell Suspension Culture

The growing of cells including the culture of single cells or small

TN_GOVT_BIOLOGY-BOTANY_XII_PAGES_107-121 CH 05.indd 100 3/14/2020 10:36:55 AM