and reproducing ability heterosis can be classified as:

i. **Euheterosis**- This is the true heterosis which is inherited and is further classified as:

a. **Mutational Euheterosis** - Simplest type of euheterosis and results from the sheltering or eliminating of the deleterious, unfavourable often lethal, recessive, mutant genes by their adaptively superior dominant alleles in cross pollinated crops.

b. **Balanced Euheterosis** – Well balanced gene combination which is more adaptive to environmental conditions and agricultural usefulness.

ii. **Pseudoheterosis** – Also termed as luxuriance. Progeny possess superiority over parents in vegetative growth but not in yield and adaptation, usually sterile or poorly fertile.

9.7.5. Mutation Breeding

Muller and Stadler (1927- 1928) coined the term mutation breeding. It represents a new method of conventional breeding procedures as they have the advantage of improving the defect without losing agronomic and quality character in agriculture and crop improvement. Mutation means the sudden heritable changes in the genotype or phenotype of an organism. Gene mutations are of considerable importance in plant breeding as they provide essential inputs for evolution as well as for re-combination and selection. It is the only method for improving seedless crops.

Radiation such as UV short wave, X-ray, Alpha (α), Beta (β), Gamma waves and many chemicals such as cesium, EMS (ethyl methane sulfonate), nitromethylurea induce mutation to develop new varieties of crops. **Example:** Triple gene dwarf wheat with increase in yield and height. Atomita 2 - rice with salinity tolerance and pest resistance.

9.7.6. Polyploid Breeding

Majority of flowering plants are diploid (2n). The plants which possess more than two sets of chromosome are called polyploids. Polyploidy is a major force in the evolution of both wild and cultivated plants. Polyploidy often exhibits increased hybrid vigour, increased heterozygosity, increase tolerance to both biotic and abiotic stresses, buffering of deleterious mutations. In addition, polyploidy often results in reduced fertility due to meiotic error allowing the production of seedless varieties.



Gamma Garden or Atomic Garden: Is a form of mutation breeding where plants are exposed to radioactive sources typically cobalt-60 or caesium-137 in order to generate desirable mutation in crop plants. The first Gamma garden in India is Bose Research Institute at Calcutta in 1959 and the second is IARI in 1960 which produced large variation in short type.

When chromosome number is doubled by itself in the same plant, is called autopolyploidy. Example: A triploid condition in sugarbeets, apples and pear has resulted in the increase in vigour and fruit size, large root size, large leaves, flower, more seeds and sugar content in them. It also resulted in seedless tomato, apple, watermelon and orange. Polyploidy can be induced by the use of colchicine to double the chromosome number. Allopolyploids are produced by multiplication of chromosome sets that are initially derived from two different species. Example: *Triticale (Triticum durum x secale cereale) Raphanobrassica (Brassica oleraceae x Raphanus sativus)*.

9.7.7. Green Revolution

Green revolution the term was coined by William S.Gaud in (1968). It is defined as the cumulative result of a series of research, development, innovation and technology transfer initiatives. Agricultural production (especially wheat and rice) manifolds worldwide particularly in the developing countries between the 1940's and the late 1960's.

The Green revolution or third Agricultural Revolution is the intensive plan of 1960's to increase crop yield in developing countries by introducing the high yielding, resistant varieties, increased irrigation facilities, fertilizer application and better agricultural management.

TN_GOVT_BIOLOGY-BOTANY_XII_PAGES_185-199 CH 09.indd 178 3/14/2020 11:49:13 AM