7. Functional megaspore in angiosperm develops into an (NEET 2017)
a) endosperm b) **Embryo sac**
c) embryo d) ovule

8. Which of the statement is not true. (NEET 2016)
a) Pollen grain of many species cause severe allergies
b) Stored pollen in liquid nitrogen can be used in crop breeding programmes
c) **Tapetum helps in the dehiscence of anther**
d) Exine of pollen grains is made up of sporopollenin

9) When a diploid female plant is crossed with a tetraploid male, the ploidy of endosperm cells in the resulting seed is (AIPMT 2004)
a) pentaploidy b) diploidy
c) triploidy d) **tetraploidy**

10) Which one of the following pairs of plant structures has haploid number of chromosomes? (AIPMT 2008)
a) Egg nucleus and secondary nucleus
b) Megaspore mother cell and antipodal cells
c) **Egg cell and and antipodal cells**
d) Nucellus and antipodal cells

11) The arrangement of nuclei in a normal embryo sac in the dicot plant is (AIPMT 2006)
a) 2 + 4 + 2 b) **3 + 2 + 3**
c) 2 + 3 + 3 d) 3 + 3 + 2

12) Wind pollinated flowers are (AIPMT PRE 2010)
a) Small, producing nectar and dry pollen
b) small, brightly colored, producing large number of pollen grains
c) **small, producing large number of pollen grains**
d) large, producing abundant nectar and pollen

13) Function of filiform apparatus is to (AIPMT 2014)
a) recognize the suitable pollen at stigma
b) stimulate division of generative cell
c) produce nectar
d) **guide the entry of pollen tube**

14) The coconut water from tender coconut represents (NEET 2016)
a) endocarp
b) fleshy mesocarp
c) free nuclear proembryo
d) **free nuclear endosperm**

15) Pollination in water hyacinth and water lily is brought about by the agency of (NEET 2016)
a) **insects or wind** b) birds
c) bats d) water

16) Perisperm differs from endosperm in (NEET 2013)
a) being haploid tissue
b) having no reserve food
c) **being a diploid tissue**
d) its formation by fusion of secondary nucleus with several sperms

17) Male gametes in angiosperms are formed by the division of (AIPMT 2007)
a) microspore mother cell b) microspore
c) **generative cell** d) vegetative cell

18) In a type of apomixes known as adventive polyembryony,embryo develop directly from the (AIPMT 2005)
a) synergids or antipodals in an embryo sac
b) **nucellus or integuments**
c) zygote
d) accessory embryo sac in the ovule

19) In a cereal grain the single cotyledon of the embryo is represented by (AIPMT 2006)
a) coleorhizae b) **scutellum**
c) prophyll d) coleoptiles

TN_GOVT_BIOLOGY-BOTANY_XII_PAGES_200-238 CH 10.indd 210 3/14/2020 12:18:02 PM