23. The rate of formation of new organic matter by rabbit in a grassland is called (Mains 2012)
a) net productivity
b) **secondary productivity**
c) net primary productivity
d) gross primary productivity

24. The second stage of hydrosere is occupied by plants like (Mains 2012)
a) *Azolla* b) *Typha*
c) *Salix* d) ***Vallisneria***

25. Which one of the following is a characteristic feature of cropland ecosystem? (NEET 2016)
a) Ecological succession
b) Absence of soil organisms
c) **Least genetic diversity**
d) Absence of weeds

26. Most animals that live in deep oceanic waters are (Re-AIPMT 2015)
a) **Detritivores**
b) Primary consumers
c) Secondary consumers
d) Tertiary consumers

27. During ecological succession (Re-AIPMT 2015)
a) The changes lead to a community that is in near equilibrium with the environment and is called pioneer community.
b) **The gradual and predictable change in species composition occurs in a given area**.
c) The establishment of a new biotic community is very fast in its primary phase.
d) The number and types of animals remain constant.


28. The mass of living material at a trophic level at a particular time is called (AIPMT 2015)
a) **Standing crop**
b) Gross primary productivity
c) Standing state
d) Net primary productivity

29. Match the following and select the **correct** option (AIPMT 2014)

Column I	**Column II**
(I) Earthworm	(i) pioneer species
(II) Succession	(ii) Detritivore
(III) Ecosystem service	(iii) Natality
(IV) Population growth	(iv) Pollination

	I	II	III	IV
a)	i	ii	iii	iv
b)	iv	i	iii	ii
c)	iii	ii	iv	i
d)	**ii**	**i**	**iv**	**iii**

30. Given below is a simplified model of phosphorous cycling in a terrestrial ecosystem with four blanks (A – D. Identify the blanks. (AIPMT 2014)



	A	B	C	D
a)	Rock minerals	Detritus	Litter fall	Producers
b)	Litter fall	Producers	Rock minerals	Detritus
c)	**Detritus**	**Rock minerals**	**Producers**	**Litter fall**
d)	Producers	Litter fall	Rock minerals	Detritus

31. If 20 J of energy is trapped at producer level, then how much energy will be available to peacock as food in the following chain? (AIPMT 2014)
Plant → Mice → Snake → Peacock
a) **0.02 J** b) 0.002 J
c) 0.2 J d) 0.0002 J

TN_GOVT_BIOLOGY-BOTANY_XII_PAGES_200-238 CH 10.indd 219 3/14/2020 12:18:02 PM