Diagnostic Features

- Ovule or megasporangium is protected by one / two coverings called integuments.
- The stalk of the ovule is called funicle.
- The point of attachment of funicle to the body of the ovule is known as hilum.
- The body of the ovule is made up of a central mass of parenchymatous tissue called nucellus.
- The integuments form a pore called micropyle and the region opposite to the micropyle is called as chalaza.
- The nucellus has a large, oval, sac like structure towards the micropylar end called embryo sac.
- A mature ovule, has 8 nuclei in its embryo sac.




Figure 2: L.S of ovule

Exercise 3: T.S. of *Nerium* Leaf

Aim: To observe and understand the xerophytic adaptations found in *Nerium* leaves for living in dry or xeric habitat.

Principle: The plants which are living in dry or xeric condition are known as Xerophytes.

Requirements: *Nerium* leaf, few pieces of carrot / pith / styrofoam, blade, brush, needle, compound microscope, glycerine, coverslip, wash glass, microslide, saffranin solution, petri dish, etc.



Start cutting transverse sections of *Nerium* leaf placing it in between a piece of carrot. Select the thinnest section of the material with the help of a delicate brush. Take a clean watch glass with water, transfer thin sections of the material. Put a few drops of safranin stain in the watch glass with water. Leave it for 3-5 minutes. Drain off stain and wash with water if necessary. Put the thinnest section in the centre of the slide. Put a drop of glycerine over the material. Cover it with a coverslip with the help of needle. Observe it under a compound microscope.




Figure 3: T.S. of Nerium leaf

TN_GOVT_XII_BIOLOGY_BOTANY_PRACTICAL_MANUAL_PAGES_239-258.indd 227 3/14/2020 12:12:58 PM