


Figure 3.24 Deletion




Figure 3.25 Duplication

Summary

Chromosomal theory of inheritance states that Mendelian factors have specific locus on chromosomes and they carry information from one generation to the next generation. Genes located close together on the same chromosome and inherited together are called linked genes the phenomenon is called Linkage. Two types of linkage are complete linkage and incomplete linkage. The groups of linearly arranged linked genes are called Linkage groups. Crossing over is a biological process that produces new combination of genes by inter-changing the corresponding segments between non-sister chromatids of homologous pair of chromosomes. In this segments of DNA are broken and recombined to produce new combinations of alleles a process is called Recombination. The diagrammatic representation of distances between the adjacent genes which is directly proportional to the frequency of recombination between them is called genetic mapping. When any of the three or more allelic forms of a gene occupy the same locus in a given pair of homologous chromosomes, they are said to be multiple alleles. They are m, M1 and M2 of a single gene. Mutational events that take place within individual genes are called gene mutations or point mutation, whereas the changes that occur in structure and number of chromosomes are called chromosomal mutation. The agents which are responsible for mutation is called mutagens.

Evaluation

B3WQYL

1. An allohexaploidy contains
   a) Six different genomes
   b) Six copies of three different genomes
   c) Two copies of three different genomes
   d) Six copies of one genome
2. The A and B genes are 10 cM apart on a chromosome. If an AB/ab heterozygote is testcrossed to ab/ab, how many of each progeny class would you expect out of 100 total progeny?
   a) 25 AB, 25 ab, 25 Ab, 25 aB
   b) 10 AB, 10 ab
   c) 45 AB, 45 ab
   d) 45 AB, 45 ab, 5 Ab, 5aB

TN_GOVT_BIOLOGY-BOTANY_XII_PAGES_053-077 CH 03.indd 65 3/14/2020 10:32:16 AM