and grow, can accept a range of vectors and has also been studied for safety. One more important feature of *E.coli* to be preferred as a host cell is that under optimal growing conditions the cells divide every 20 minutes.

Since the DNA is a hydrophilic molecule,it cannot pass through cell membranes, In order to force bacteria to take up the plasmid, the bacterial cells must first be made competent to take up DNA. This is done by treating them with a specific concentration of a divalent cation such as calcium. Recombinant DNA can then be forced into such cells by incubating the cells with recombinant DNA on ice, followed by placing them briefly at 42^{0}C (heatshock) and then putting them back on ice. This enables bacteria to take up the Recombinant DNA.

For the expression of eukaryotic proteins, eukaryotic cells are preferred because to produce a functionally active protein it should fold properly and post translational modifications should also occur, which is not possible by prokaryotic cell (E.coli).




Figure 4.15: Liposome mediated method of Gene Transfer

4.6 Methods of Gene Transfer

The next step after a recombinant DNA molecule has been generated is to introduce it into a suitable host cell. There are many methods to introduce recombinant vectors and these are dependent on several factors such as the vector type and host cell.

For achieving genetic transformation in plants, the basic pre-requisite is the construction of a vector which carries the gene of interest flanked by the necessary controlling sequences, i.e., the promoter and terminator, and deliver the genes into the host plant. There are two kinds of gene transfer methods in plants. It includes:

- Direct or vectorless gene transfer
- Indirect or vector – mediated gene transfer

4.6.1 Direct or Vectorless Gene Transfer

In the direct gene transfer methods, the foreign gene of interest is delivered into the host plant without the help of a vector. The following are some of the common methods of direct gene transfer in plants.

a. **Chemical mediated gene transfer:** Certain chemicals like polyethylene glycol (PEG) and dextran sulphate induce DNA uptake into plant protoplasts.

b. **Microinjection:** The DNA is directly injected into the nucleus using fine tipped glass needle or micro pipette to transform plant cells. The protoplasts are immobilised on a solid support (agarose on a microscopic slide) or held with a holding pipette under suction.

c. **Electroporation Methods of Gene Transfer:** A pulse of high voltage is applied to protoplasts, cells or tissues which makes transient pores in the plasma membrane through which uptake of foreign DNA occurs.




Figure 4.14: Electroporation Methods of Gene Transfer

TN_GOVT_BIOLOGY-BOTANY_XII_PAGES_078-106 CH 04.indd 76 3/14/2020 10:33:51 AM