and r be the radius of the oil drop, then the mass of the oil drop can be expressed in terms of its density as

$$\rho = \frac{m}{V}$$

$$\Rightarrow m = \rho\left(\frac{4}{3}\pi r^3\right) \qquad \left(\because \text{volume of the sphere}, V = \frac{4}{3}\pi r^3\right)$$

The gravitational force can be written in terms of density as

$$F_g = mg \Rightarrow F_g = \rho\left(\frac{4}{3}\pi r^3\right)g$$

Let σ be the density of the air, the upthrust force experienced by the oil drop due to displaced air is

$$F_b = \sigma\left(\frac{4}{3}\pi r^3\right)g$$

Once the oil drop attains a terminal velocity υ, the net downward force acting on the oil drop is equal to the viscous force acting opposite to the direction of motion of the oil drop. From Stokes law, the viscous force on the oil drop is

$$F_v = 6\pi r\upsilon\eta$$

From the free body diagram as shown in Figure 9.7 (a), the force balancing equation is

$$F_g = F_b + F_\upsilon$$

$$\rho\left(\frac{4}{3}\pi r^3\right)g = \sigma\left(\frac{4}{3}\pi r^3\right)g + 6\pi r\upsilon\eta$$

$$\frac{4}{3}\pi r^3(\rho-\sigma)g = 6\pi r\upsilon\eta$$

$$\frac{2}{3}\pi r^3(\rho-\sigma)g = 3\pi r\upsilon\eta$$

$$r = \left[\frac{9\eta\upsilon}{2(\rho-\sigma)g}\right]^{\frac{1}{2}} \qquad (9.9)$$

Thus, equation (9.9) gives the radius of the oil drop.

(b) Determination of electric charge

When the electric field is switched on, charged oil drops experience an upward electric force (qE). Among many drops, one particular drop can be chosen in the field of view of microscope and strength of the electric field is adjusted to make that particular drop to be stationary. Under these circumstances, there will be no viscous force acting on the oil drop. Then, from the free body diagram shown Figure 9.7 (b), the net force acting on the oil droplet is

$$F_e + F_b = F_g$$

$$\Rightarrow qE + \frac{4}{3}\pi r^3\sigma g = \frac{4}{3}\pi r^3\rho g$$

$$\Rightarrow qE = \frac{4}{3}\pi r^3(\rho-\sigma)g \qquad (9.10)$$

$$\Rightarrow q = \frac{4}{3E}\pi r^3(\rho-\sigma)g \qquad (9.11)$$

Substituting equation (9.9) in equation (9.11), we get

$$q = \frac{18\pi}{E}\left(\frac{\eta^3\upsilon^3}{2(\rho-\sigma)g}\right)^{\frac{1}{2}} \qquad (9.12)$$

Millikan repeated this experiment several times and computed the charges on oil drops. He found that the charge of any oil drop can be written as integral multiple of a basic value, -1.6×10^{-19}C,

UNIT-9(XII-Physics_Vol-2)_New.indd 147 21-07-2020 15:48:30