Multiple choice questions

1. The speed of light in an isotropic medium depends on,

(a) its intensity

(b)its wavelength

(c) the nature of propagation

(d) the motion of the source w.r.t medium

D7F8Y5

2. A rod of length 10 cm lies along the principal axis of a concave mirror of focal length 10 cm in such a way that its end closer to the pole is 20 cm away from the mirror. The length of the image is, (AIPMT Main 2012)

(a) 2.5 cm (b) 5cm

(c) 10 cm (d) 15cm

3. An object is placed in front of a convex mirror of focal length of f and the maximum and minimum distance of an object from the mirror such that the image formed is real and magnified. (JEE Main 2009)]

(a) $2f$ and c (b) c and ∞

(c) f and O (d) None of these

4. For light incident from air on a slab of refractive index 2, the maximum possible angle of refraction is,

(a) 30° (b) 45°

(c) 60° (d) 90°

5. If the velocity and wavelength of light in air is V_a and λ_a and that in water is V_w and λ_w, then the refractive index of water is,

(a) $\frac{V_w}{V_a}$ (b) $\frac{V_a}{V_w}$

(c) $\frac{\lambda_w}{\lambda_a}$ (d) $\frac{V_a\lambda_a}{V_w\lambda_w}$

6. Stars twinkle due to,

(a) reflection

(b) total internal reflection

(c) refraction

(d) polarisation

7. When a biconvex lens of glass having refractive index 1.47 is dipped in a liquid, it acts as a plane sheet of glass. This implies that the liquid must have refractive index,

(a) less than one

(b) less than that of glass

(c) greater than that of glass

(d) equal to that of glass

8. The radius of curvature of curved surface at a thin planoconvex lens is 10 cm and the refractive index is 1.5. If the plane surface is silvered, then the focal length will be,

(a) 5 cm (b) 10 cm

(c) 15 cm (d) 20 cm

9. An air bubble in glass slab of refractive index 1.5 (near normal incidence) is 5 cm deep when viewed from one surface and 3 cm deep when viewed from the opposite face. The thickness of the slab is,

(a) 8 cm (b) 10 cm

(c) 12 cm (d) 16 cm

10. A ray of light travelling in a transparent medium of refractive index n falls, on a surface separating the medium from air at an angle of incidents of 45°. The ray can undergo total internal reflection for the following n,

(a) $n = 1.25$ (b) $n = 1.33$

(c) $n = 1.4$ (d) $n = 1.5$

UNIT-6(XII-Physics_Vol-2)_new.indd 48 21-07-2020 16:47:55