removed, or altered at particular locations in the genome. Several approaches to genome editing have been developed. A recent one is known as CRISPR-Cas9, which is short form of **Clustered Regularly Interspaced Short Palindromic Repeats** and CRISPR-associated protein 9. The CRISPR-Cas9 system has generated a lot of excitement in the scientific community because it is faster, cheaper, more accurate, and more efficient than other existing genome editing methods.

Rice, was among the first plants to be used to demonstrate the feasibility of CRISPR-mediated targeted mutagenesis and gene replacement. The gene editing tool CRISPR can be used to make hybrid rice plants that can clone their seed. Imtiyaz Khand and Venkatesan Sundaresan and colleagues reported in a new study which clearly shows one can re-engineer rice to switch it from a sexual to an asexual mode.

4.7.10 RNA Interference (RNAi)

All characters of organism are the result of expression of different genes which are regions of nuclear DNA. This expression involves transcription and translation. Transcription refers to the copying of genetic information from one strand of the DNA (called sense strand) by RNA. This RNA, as soon as it formed cannot be straight away sent to the cytoplasm to undertake the process of translation. It has to be edited and made suitable for translation which brings about protein synthesis. One of the main items removed from the RNA strand are the introns. All these changes before translation normally take place whereby certain regions of DNA are silence. However, there is an (RNAi) pathway. RNA interference is a biological process in which RNA molecules inhibit gene expression or translation. This is done by neutralising targetd mRNA molecules.

A simplified model for the RNAi pathway is based on two steps, each involving




Figure 4.23: RNA Interference

ribonuclease enzyme. In the first step, the trigger RNA (either dsRNA or miRNA primary transcript) is processed into a short interfering RNA (siRNA) by the RNase II enzymes called Dicer and Drosha. In the second step, siRNAs are loaded into the effector complex RNA-induced silencing complex (RISC). The siRNA is unwound during RISC assembly and the single-stranded RNA hybridizes with mRNA target. This RNAi is seen in plant feeding nematodes.

4.8 Transgenic Plants / Genetically Modified Crops (Gm Crops)

4.8.1 Herbicide Tolerant – Glyphosate

Weeds are a constant problem in crop fields. Weeds not only compete with crops for sunlight, water, nutrients and space but also acts as a carrier for insects and diseases. If left uncontrolled, weeds can reduce crop yields significantly.

Transgenic plants contain a novel DNA introduced into the genome.

Glyphosate herbicide produced by Monsanto, USA company under the trade name 'Round up' kills plants by blocking the 5-enopyruvate shikimate-3 phosphate synthase (EPSPS) enzyme, an enzyme involved in the biosynthesis of aromatic amino acids, vitamins and many secondary plant metabolites. There are several ways by which crops can be modified to be glyphosate-tolerant.

TN_GOVT_BOTANY_XII_PAGES_092-120 CH 04.indd 96 3/8/2020 2:45:36 PM