the atmosphere increases due to burning of fossile fuels, deforestration, forest fire, volcanic eruption and decomposition of dead organic matters. The details of carbon cycle are given in the figure.

Phosphorus cycle

It is a type of sedimentary cycle. Already we know that phosphorus is found in the biomolecules like DNA, RNA, ATP, NADP and phospholipid molecules of living organisms. Phosphorus is not abundant in the biosphere, whereas a bulk quantity of phosphorus is present in rock deposits, marine sediments and guano. It is released from these deposits by weathering

7.2.10 Types of ecosystem

Biosphere consists of different types of ecosystems, which are as follows:




Figure 7.14: Types of Ecosystem

Though there are many types of ecosystems as charted above. Only the pond ecosystem is detailed below.

Structure of Pond ecosystem




Figure 7.15: Diagram shows structure of pond ecosystem with abiotic and biotic components.

It is a classical example for natural, aquatic, freshwater, lentic type of ecosystem. It helps us to understand the structure and function of an ecosystem. When rain water gathers in a shallow area, gradually over a period of time, different kinds of organisms (microbes, plants, animals) become part of this ecosystem. This pond ecosystem is a self sustaining and self regulatory fresh water ecosystem, which shows a complex interaction between the abiotic and biotic components in it.

Activity

Collect few living and nonliving components from any water body found near by.

Abiotic components

A pond ecosystem consists of dissolved inorganic (CO_2, O_2, Ca, N, Phosphate) and organic substances (amino acids and humic acid) formed from the dead organic matter. The function of pond ecosystem is regulated by few factors like the amount of light, temperature, pH value of water and other climatic conditions.

Biotic components

They constitute the producers, variety of consumers and decomposers (microorganisms).

TN_GOVT_BOTANY_XII_PAGES_162-190 CH 07.indd 158 3/8/2020 2:54:02 PM