it will be gradually covered by plant community again and become fertile. Such successive replacement of one type of plant community by the other of the same area/ place is known as plant **succession**. The first invaded plants in a barren area are called **pioneers**. On the other hand, a series of transitional developments of plant communities one after another in a given area are called **seral communities**. At the end a final stage and a final plant community gets established which are called as climax and climax community respectively.

7.3.1 Causes of Succession

Ever since the onset of origin of life, organic evolution and ecological succession are taking place parallelly. Ecological succession is a complex process. There are three types of causes for any ecological succession. They are

a. **Initiating causes** - Activity of abiotic (light, temperature, water, fire, soil erosion and wind) and biotic factors (competition among organisms) leads to formation of a barren area or destruction of the existing community of an area, initiating primary or secondary succession respectively.

b. **Continuing causes** - The processes of migration, aggregation, competition, reaction etc, are the continuing causes which lead to change the plant communities and nature of the soil in an area.

c. **Stabilizing causes** - The stabilization of the plant community in an area is primarly controlled by climatic factors rather than other factors.

7.3.2 Characteristics of ecological succession

- It is a systematic process which causes changes in specific structure of plant community.
- It is resultant of changes of abiotic and biotic factors.
- It transforms unstable community into a stable community.
- Gradual progression in species diversity, total biomass, niche specialisation, and humus content of soil takes place.
- It progresses from simple food chain to complex food web.
- It modifies the lower and simple life form to the higher life forms.
- It creates inter-dependence of plants and animals.

7.3.3 Types of succession

The various types of succession have been classified in different ways on the basis of different aspects. These are as follows:

1. Primary succession - The development of plant community in a barren area where no community existed before is called primary succession. The plants which colonize first in a barren area is called **pioneer species** or **primary community** or **primary colonies**. Generally, Primary succession takes a very long time for the occurrence in any region.

Example: Microbes, Lichen, Mosses.

2. Secondary succession - The development of a plant community in an area where an already developed community has been destroyed by some natural disturbance (Fire, flood, human activity) is known as **secondary succession**. Generally, This succession takes less time than the time taken for primary succession.

Example: The forest destroyed by fire and excessive lumbering may be re-occupied by herbs over a period of time.




Figure 7.18: Diagrammatic representation of secondary succession

TN_GOVT_BOTANY_XII_PAGES_162-190 CH 07.indd 163 3/8/2020 2:54:04 PM