


Figure 1.6 Development of male gametophyte

1.4.2 Female reproductive part - Gynoecium

The **gynoecium** represents the female reproductive part of the flower. The word gynoecium represents one or more pistils of a flower. The word pistil refers to the ovary, style and stigma. A pistil is derived from a carpel. The word ovary represents the part that contains the ovules. The stigma serves as a landing platform for pollen grains. The style is an elongated slender part beneath the stigma. The basal swollen part of the pistil is the ovary. The ovules are present inside the ovary cavity (locule)on the placenta .Gynoecium (carpel) arises as a small papillate outgrowth of meristematic tissue from the growing tip of the floral primordium. It grows actively and soon gets differentiated into ovary, style and stigma. The ovules or megasporangia arise from the placenta. The number of ovules in an ovary may be one (paddy, wheat and mango) or many (papaya, water melon and orchids).

Structure of ovule(Megasporangium):

Ovule is also called megasporangium and is protected by one or two covering called **integuments**. A mature ovule consists of a stalk and a body. The stalk or the **funiculus** (also called funicle) is present at the base and it attaches the ovule to the placenta. The point of attachment of funicle to the body of the ovule is known as **hilum**. It represents the junction between ovule and funicle. In an inverted ovule, the funicle is adnate to the body of the ovule forming a ridge called **raphe**. The body of the ovule is made up of a central mass of parenchymatous tissue called **nucellus** which has large reserve food materials. The nucellus is enveloped by one or two protective coverings called **integuments**. Integument encloses the nucellus completely except at the top where it is free and forms a pore called micropyle. The ovule with one or two integuments are said to be **unitegmic** or **bitegmic** ovules respectively. The basal region of the body of the ovule where the nucellus, the integument and the funicle meet or merge is called as **chalaza**. There is a large,




Figure 1.7 Structure of an ovule

oval, sac-like structure in the nucellus toward the micropylar end called **embryo sac** or female gametophyte. It develops from the functional megaspore formed within the nucellus. In some species(unitegmic tenuinucellate) the inner layer of the integument may become specialized to perform the nutritive function for the embryo

TN_GOVT_BOTANY_XII_PAGES_001-034 CH 01.indd 10 3/8/2020 2:38:42 PM