megaspores are usually arranged in a linear tetrad. Of the four megaspores formed, usually the chalazal one is functional and other three megaspores degenerate. The functional megaspore forms the female gametophyte or embryo sac. This type of development is called **monosporic** development (Example: *Polygonum*). Of the four megaspores formed if two are involved in Embryo sac formation the development is called **bisporic** (Example: *Allium*). If all the four megaspores are involved in Embryo sac formation the development is called **tetrasporic** (Example: *Peperomia*). An ovule generally has a single embryo sac. The development of monosporic embryo sac (*Polygonum* type) is given in Figure 1.9.

Development of Monosporic embryo sac.

To describe the stages in embryo sac development and organization the simplest monosporic type of development is given below.

The functional megaspore is the first cell of the embryo sac or female gametophyte. The megaspore elongates along micropylar- chalazal axis. The nucleus undergoes a mitotic division. Wall formation does not follow the nuclear division. A large central vacuole now appears between the two daughter nuclei. The vacuole expands and pushes the nuclei towards the




Figure 1.10 Structure of Embryo sac

opposite poles of the embryo sac. Both the nuclei divide twice mitotically, forming four nuclei at each pole. At this stage all the eight nuclei are present in a common cytoplasm (free nuclear division). After the last nuclear division the cell undergoes appreciable elongation, assuming a sac-like appearance. This is followed by cellular organization of the embryo sac. Of the four nuclei at the micropylar end of the embryo sac, three organize into an **egg apparatus**, the fourth one is left free in the cytoplasm of the central cell as the upper polar nucleus. Three nuclei of the chalazal end form three **antipodal cells** whereas the fourth one functions as the lower polar nucleus. Depending on the plant the **2 polar nuclei** may remain free or may fuse to form a **secondary nucleus** (central cell). The egg apparatus is made up of a central egg cell and two synergids, one on each side of the egg cell. Synergids secrete chemotropic substances that help to attract the pollen tube. The special




Figure 1.9 Development of ovule and embryo sac (*Polygonum type*).

TN_GOVT_BOTANY_XII_PAGES_001-034 CH 01.indd 12 3/8/2020 2:38:43 PM