13) Function of filiform apparatus is to (AIPMT 2014)
a) recognize the suitable pollen at stigma
b) stimulate division of generative cell
c) produce nectar
d) **guide the entry of pollen tube**

14) The coconut water from tender coconut represents (NEET 2016)
a) endocarp
b) fleshy mesocarp
c) free nuclear proembryo
d) **free nuclear endosperm**

15) Pollination in water hyacinth and water lily is brought about by the agency of (NEET 2016)
a) **insects or wind** b) birds
c) bats d) water

16) Perisperm differs from endosperm in (NEET 2013)
a) being haploid tissue
b) having no reserve food
c) **being a diploid tissue**
d) its formation by fusion of secondary nucleus with several sperms

17) Male gametes in angiosperms are formed by the division of (AIPMT 2007)
a) microspore mother cell b) microspore
c) **generative cell** d) vegetative cell

18) In a type of apomixes known as adventive polyembryony,embryo develop directly from the (AIPMT 2005)
a) synergids or antipodals in an embryo sac
b) **nucellus or integuments**
c) zygote
d) accessory embryo sac in the ovule

19) In a cereal grain the single cotyledon of the embryo is represented by (AIPMT 2006)
a) coleorhizae b) **scutellum**
c) prophyll d) coleoptiles

20) An ovule which becomes curved so that the nucellus and embryo sac lie at right angles to the funicle is (AIPMT 2004)
a) camylotropous b) anatropous
c) orthotropous d) **hemianatropous**

21) Endosperm is formed during the double fertilization by (AIPMT 2000)
a) **two polar nuclei and one male gamete**
b) one polar nuclei and one male gamete
c) ovum and male gametes
d) two polar nuclei and two male gametes

UNIT VII – Genetics

1. Genes for cytoplasmic male sterility in plants are generally located in (AIPMT 2005)
a) **Mitrochondrial genome** b) Cytosol
c) Chloroplast genome d) Nuclear genome

2. In which mode of inheritance do you expect more maternal influence among the off spring (AIPMT 2006)
a) Autosomal b) **Cytoplasmic**
c) Y-linked d) X-linked

3. Which one of the following cannot be explained on the basis of Mendel's Law of Dominance? (AIPMT 2010)
a) Factors occur in pairs
b) The discrete unit controlling a particular character is called a factor
c) Out of one pair of factors one is dominant and the other is recessive
d) Alleles does not show any blending and both the characters recover as such in F_2 generation

4. F_2 generation in a Mendelian cross shows that both genotypic and phenotypic ratios are same as 1:2:1. It represents a case of (AIPMT 2012)
a) Monohybrid crosses with incomplete dominance
b) Co-dominance c) Dihybrid cross
d) Monohybrid cross with complete dominance

TN_GOVT_BOTANY_XII_PAGES_230-283 CH 10.indd 256 3/8/2020 2:59:37 PM