

a) Level one PP is 'pipal trees' and the level SC is 'sheep'.

b) Level PC is 'rats' and level SC is 'cats'

c) **Level PC is 'insects' and level SC is 'small insectivorous birds'**

d) Level PP is 'phytoplanktons' in sea and 'whale' on top level TC

33. Which one of the following statements for pyramid of energy is incorrect, whereas the remaining three are correct? (AIPMT Prelims 2011)

a) It is upright in shape

b) Its base is broad

c) It shows energy content of different trophic level organisms

d) **It is inverted in shape**

34. Which one of the following animals may occupy more than one trophic levels in the same ecosystem at the same time? (AIPMT Mains 2011)

a) Goat b) Frog

c) **Sparrow** d) Lion

35. Both hydrarch and xerarch successions lead to (AIPMT Mains 2011)

a) Highly dry conditions

b) Excessive wet conditions

c) **Medium water conditions**

d) Xeric conditions

36. Of the total incident solar radiation the proportion of PAR is (AIPMT Prelims 2011)

a) More than 80% b) About 70%

c) About 60% d) **Less than 50%**

37. The breakdown of detritus into smaller particles by earthworm is a process called (AIPMT Mains 2011)

a) Mineralisation b) Catabolism

c) Humification d) **Fragmentation**

38. The biomass available for consumption by the herbivores and the decomposers is called (AIPMT Prelims 2010)

a) Gross primary productivity

b) **Net primary productivity**

c) Secondary productivity

d) Standing crop

39. The correct sequence of plants in a hydrosere is (AIPMT Prelims 2009)

a) ***Volvox*** → ***Hydrilla*** → ***Pistia*** → ***Scirpus*** → ***Lantana*** → **Oak**

b) *Pistia*→*Volvox*→*Scirpus*→*Hydrilla*→Oak → *Lantana*

c) Oak→*Lantana*→*Volvox*→*Hydrilla*→*Pistia* →*Scirpus*

d) Oak→*Lantana*→*Scirpus*→*Pistia*→*Hydrilla* →*Volvox*

40. About 70% of the total global carbon is found in (AIPMT Prelims 2008)

a) Forests b) Grasslands

c) Agro ecosystems d) **Oceans**

41. Consider the following statements concerning food chains

i) Removal of 80% tigers from an area resulted in greatly increased growth of vegetation.

ii) Removal of most of the carnivores resulted in an increased population of deers.

iii) The length of food chains is generally limited to 3 – 4 trophic levels due to energy loss.

iv) The length of food chains may vary from 2 to 8 trophic levels.

Which two of the above statements are correct? (AIPMT Prelims 2008)

a) i and ii b) **ii and iii**

c) iii and iv d) i and iv

TN_GOVT_BOTANY_XII_PAGES_230-283 CH 10.indd 267 3/8/2020 2:59:37 PM