Bulbils – *Fritillaria imperialis*; Bulbs – *Allium*; Runner – *Mentha arvensis*; Sucker - *Chrysanthemum*

Agamospermy: It refers to processes by which Embryos are formed by eliminating meiosis and syngamy.

Adventive embryony

An Embryo arises directly from the diploid sporophytic cells either from nucellus or integument. It is also called **sporophytic budding** because gametophytic phase is completely absent. Adventive embryos are found in *Citrus* and *Mangifera*

Diplospory (Generative apospory): A diploid embryo sac is formed from megaspore mother cell without a regular meiotic division Examples. *Eupatorium* and *Aerva.*

Apospory: Megaspore mother cell (MMC) undergoes the normal meiosis and four megaspores formed gradually disappear. A nucellar cell becomes activated and develops into a diploid embryo sac. This type of apospory is also called somatic apospory. Examples *Hieracium* and *Parthenium.*

1.8 Polyembryony

Occurrence of more than one embryo in a seed is called polyembryony (Figure 1.24). The first case of polyembryony was reported in certain oranges by Anton von Leeuwenhoek in the year 1719. Polyembryony is divided into four categories based on its origin.

a. **Cleavage polyembryony** (Example: Orchids)

b. **Formation of embryo by cells of the Embryo sac other than egg** (Synergids – *Aristolochia*; antipodals – *Ulmus* and endosperm – *Balanophora*)

c. **Development of more than one Embryo sac within the same ovule.** (Derivatives of same MMC, derivatives of two or more MMC – *Casuarina*)

d. **Activation of some sporophytic cells of the ovule** (Nucellus/ integuments-*Citrus and Syzygium*).




Figure 1.24 : Polyembryony – Embryo sac of *Ulmus glabra* showing zygotic and antipodal embryo

Practical applications

The seedlings formed from the nucellar tissue in *Citrus* are found better clones for Orchards. Embryos derived through polyembryony are found virus free.

1.9 Parthenocarpy

As mentioned earlier, the ovary becomes the fruit and the ovule becomes the seed after fertilization. However in a number of cases, fruit like structures may develop from the ovary without the act of fertilization. Such fruits are called **parthenocarpic fruits**. Invariably they will not have true seeds. Many commercial fruits are made seedless. Examples: Banana, Grapes and Papaya.

Significance

- The seedless fruits have great significance in horticulture.
- The seedless fruits have great commercial importance.
- Seedless fruits are useful for the preparation of jams, jellies, sauces, fruit drinks etc.
- High proportion of edible part is available in parthenocarpic fruits due to the absence of seeds.

TN_GOVT_BOTANY_XII_PAGES_001-034 CH 01.indd 27 3/8/2020 2:39:03 PM