1) Incomplete dominance (2) Codominance (3) Multiple alleles (4) Pleiotropic genes are common examples for intragenic interaction.

2.4.1. Incomplete dominance – No blending of genes

The German Botanist Carl Correns's (1905) Experiment - In 4 O' clock plant, *Mirabilis jalapa* when the pure breeding homozygous red (R^1R^1) parent is crossed with homozygous white (R^2R^2), the phenotype of the F_1 hybrid is heterozygous pink (R^1R^2). The F_1 heterozygous phenotype differs from both the parental homozygous phenotype. This cross did not exhibit the character of the dominant parent but an intermediate colour pink. When one allele is not completely dominant to another allele it shows incomplete dominance. Such allelic interaction is known as incomplete dominance. F_1 generation produces intermediate phenotype pink coloured flower. When pink coloured plants of F_1 generation were interbred in F_2 both phenotypic and genotypic ratios were found to be identical as 1 : 2 : 1(1 red : 2 pink : 1 white). Genotypic ratio is 1 R^1R^1 : 2 R^1R^2 : 1 R^2R^2.From this we conclude that the alleles themselves remain discrete and unaltered proving the Mendel's Law of Segregation. The phenotypic and genotypic ratios are the same. There is no blending of genes. In the F_2 generation R^1 and R^2 genes segregate and recombine to produce red, pink and white in the ratio of 1 : 2 : 1. R^1 allele codes for an enzyme responsible for the formation of red pigment. R^2 allele codes for defective enzyme. R^1 and R^2 genotypes produce only enough red pigments to make the flower pink. Two R^1R^1 are needed for producing red flowers. Two R^2R^2 genes are needed for white flowers. If blending had taken place, the original pure traits would not have appeared and all F_2 plants would have

How are we going to interpret the lack of dominance and give explanation to the intermediate heterozygote phenotype?

How will you explain incomplete dominance at the molecular level?

Gene expression is explained in a quantitative way. Wild-type allele which is a functional allele when present in two copies (R^1 R^1) produces an functional enzyme which synthesizes red pigments. The mutant allele which is a defective allele in two copies (R^2 R^2) produces an enzyme which cannot synthesize necessary red pigments. The white flower is due to the mutation causing complete loss of function. The F_1 intermediate phenotype heterozygote (R^1R^2) has one copy of the allele R^1. R^1 produces 50% of the functional protein resulting in half of the pigment of red flowered plant and so it is pink. The intermediate phenotype pink heterogyzote with 50% of functional protein is not enough to create the red phenotype homozygous, which makes 100% of the functional protein.

2.4.2. Codominance (1 : 2 : 1)

This pattern occurs due to simultaneous (joint) expression of both alleles in the heterozygote - The phenomenon in which two alleles are both expressed in the heterozygous individual is known as codominance. Example: Red and white flowers of *Camellia*, inheritance of sickle cell haemoglobin, ABO blood group system in humanbeings. In humanbeings, I^A and I^B alleles of I gene are codominant which follows Mendels law of segregation. The codominance was demonstrated in plants with the help of electrophoresis or chromatography for protein or flavonoid substance. Example: *Gossypium hirsutum* and *Gossypium sturtianum*, their F_1 hybrid (amphiploid) was tested for seed proteins by electrophoresis. Both the parents have different banding patterns for their seed proteins. In hybrids, additive banding pattern was noticed.

TN_GOVT_BOTANY_XII_PAGES_035-054 CH 02.indd 43 3/8/2020 2:42:39 PM