3.2.1 Coupling and Repulsion theory

The two dominant alleles or recessive alleles occur in the same homologous chromosomes, tend to inherit together into same gamete are called **coupling or *cis* configuration** (Figure: 3.5). If dominant or recessive alleles are present on two different, but homologous chromosomes they inherit apart into different gamete are called **repulsion or *trans* configuration** (Figure: 3. 6)**.**




Figure 3.4: *Cis-Trans* arrangement of genes

3.2.2 Kinds of Linkage

T.H. Morgan found two types of linkage. They are complete linkage and incomplete linkage depending upon the absence or presence of new combination of linked genes.

Complete Linkage

If the chances of separation of two linked genes are not possible those genes always remain together as a result, only parental combinations are observed. The linked genes are located very close together on the same chromosome such genes do not exhibit crossing over. This phenomenon is called **complete linkage**. It is rare but has been reported in male ***Drosophila*. C.B Bridges** (1919) discovered that crossing over is completely absent in some species of male ***Drosophila***.

Incomplete Linkage

If two linked genes are sufficiently apart, the chances of their separation are possible. As a result, parental and non-parental combinations are observed. The linked genes exhibit some crossing over. This phenomenon is called **incomplete linkage**. This was observed in maize. It was reported by Hutchinson.




Figure 3.5: Alleles in coupling or cis configuration

3.2.3 Linkage Groups

The groups of linearly arranged linked genes on a chromosome are called **Linkage groups**. In any species the number of linkage groups corresponds to the number haploid set of chromosomes. Example:

TN_GOVT_BOTANY_XII_PAGES_055-091 CH 03.indd 55 3/8/2020 2:44:12 PM