


Figure 3.6: Alleles in repulsion or trans configuration

Name of organism	Linkage groups
Mucor	2
Drosophila	4
Sweet pea	7
Neurospora	7
Maize	10

Table 3.3 : Linkage groups in some organisms

Linkage and crossing over are two processes that have opposite effects. Linkage keeps particular genes together but crossing over mixes them. The differences are given below.

Linkage	Crossing over
1. The genes present on chromosome stay close together	It leads to separation of linked genes
2. It involves same chromosome of homologous chromosome	It involves exchange of segments between non-sister chromatids of homologous chromosome.
3. It reduces new gene combinations	It increases variability by forming new gene combinations. lead to formation of new organism

Table 3.4: Differences between linkage and crossing over

3.3 Crossing Over

Crossing over is a biological process that produces new combination of genes by inter-changing the corresponding segments between non-sister chromatids of homologous pair of chromosomes. The term 'crossing over' was coined by **Morgan (1912).** It takes place during pachytene stage of prophase I of meiosis. Usually crossing over occurs in germinal cells during gametogenesis. It is called meiotic or germinal crossing over. It has universal occurrence and has great significance. Rarely, crossing over occurs in somatic cells during mitosis. It is called somatic or mitotic crossing over.

3.3.1 Mechanism of Crossing Over

Crossing over is a precise process that includes stages like synapsis, tetrad formation, cross over and terminalization.

(i) Synapsis

Intimate pairing between two homologous chromosomes is initiated during zygotene stage of prophase I of meiosis I. Homologous chromosomes are aligned side by side resulting in a pair of homologous chromosomes called **bivalents**. This pairing phenomenon is called **synapsis or syndesis.** It is of three types,

TN_GOVT_BOTANY_XII_PAGES_055-091 CH 03.indd 56 3/8/2020 2:44:12 PM