1. **Procentric synapsis:** Pairing starts from middle of the chromosome.
2. **Proterminal synapsis:** Pairing starts from the telomeres.
3. **Random synapsis:** Pairing may start from anywhere.

(ii) Tetrad Formation

Each homologous chromosome of a bivalent begin to form two identical sister chromatids, which remain held together by a centromere. At this stage each bivalent has four chromatids. This stage is called **tetrad stage**.

(iii) Cross Over

After tetrad formation, crossing over occurs in pachytene stage. The non-sister chromatids of homologous pair make a contact at one or more points. These points of contact between non-sister chromatids of homologous chromosomes are called **Chiasmata** (singular-Chiasma). At chiasma, cross-shaped or X-shaped structures are formed, where breaking and rejoining of two chromatids occur. This results in reciprocal exchange of equal and corresponding segments between them.




Figure 3.9: Structure of Synaptonemal Complex

(iv) Terminalisation

After crossing over, chiasma starts to move towards the terminal end of chromatids. This is known as **terminalisation**. As a result, complete separation of homologous chromosomes occurs. (Figure 4.10)

3.3.2 Importance of Crossing Over

Crossing over occurs in all organisms like bacteria, yeast, fungi, higher plants and animals. Its importance is

- Exchange of segments leads to new gene combinations which plays an important role in evolution.
- Studies of crossing over reveal that genes are arranged linearly on the chromosomes.
- Genetic maps are made based on the frequency of crossing over.
- Crossing over helps to understand the nature and mechanism of gene action.
- If a useful new combination is formed it can be used in plant breeding.




Figure 3.10: Mechanism of crossing over

3.3.3 Recombination

Crossing over results in the formation of new combination of characters in an organism called recombinants. In this, segments of DNA are broken and recombined to produce new combinations of alleles. This process is called **Recombination.**

Calculation of Recombination Frequency (RF)

The percentage of recombinant progeny in a cross is called recombination frequency. The recombination frequency (cross over

TN_GOVT_BOTANY_XII_PAGES_055-091 CH 03.indd 57 3/8/2020 2:44:13 PM