Recombination: In cells the genetic information within and among DNA molecule are re-arranged by a process called genetic recombination. Recombination is the result of crossing over between the pairs of homologous chromosomes during meiosis. In earlier classes you have learnt chromosomal recombination. In molecular level it involves breakage and reunion of polynucleotides.

3.7.1 Eukaryotic DNA replication

Replication starts at a specific site on a DNA sequence known as the Origin of replication. There are more than one origin of replication in eukaryotes. ***Saccharomyces cerevisiae*** (yeast) has approximately 400 origins of replication. DNA replication in eukaryotes starts with the assembly of a prereplication complex (preRC) consisting of 14 different proteins. Part of a preRC is a group of 6 proteins called the origin recognition complex (ORC) which acts as initiator in eukaryotic DNA replication. The origin of replication in **yeast** is called as **ARS sites (Autonomously Replicating Sequences).** In **yeast**, ORC was identified as a protein complex which binds directly to ARS elements.




Figure: 3.28 Eukaryotic replication fork

Replication fork is the site (point of unwinding) of separation of parental DNA strands where new daughter strands are formed. Multiple replication forks are found in eukaryotes. The enzyme **helicases** are involved in unwinding of DNA by breaking hydrogen bonds holding the two strands of DNA and replication protein A (RPA) prevents the separated polynucleotide strand from getting reattached.

Topoisomerase is an enzyme which breaks DNAs covalent bonds and removes positive supercoiling ahead of replication fork. It eliminates the torsional stress caused by unwinding of DNA double helix.

DNA replication is initiated by an enzyme **DNA polymerase α** / primase which synthesizes short stretch of RNA primers on both leading strand (continuous DNA strand) and lagging strands (discontinuous DNA strand). Primers are needed because DNA polymerase requires a free 3' OH to initiate synthesis. DNA polymerase covalently connects the nucleotides at the growing end of the new DNA strand.

DNA Pol α (alpha), DNA Pol δ (delta) and DNA Pol ε (Epsilon) are the 3 enzymes involved in nuclear DNA replication.


DNA Pol α – Synthesizes short primers of RNA

DNA Pol δ – Main Replicating enzyme of cell nucleus

DNA Pol ε – Extend the DNA Strands in replication fork

DNA Polymerase β does not play any role in the replication of normal DNA. Function - Removing incorrect bases from damaged DNA. It is involved in Base excision repair.

DNA Synthesis takes place in 5' → 3' direction and it is semidiscontinuous. When DNA is synthesized in 5' → 3' direction, only in the free 3' end (OH end) DNA is elongated. In 1960s Reiji Okazaki and his colleagues found out that one of the new DNA strands is synthesized in short pieces called **Okazaki fragments**. In discontinuous strand where the Okazaki fragments are united by ligase is called



TN_GOVT_BOTANY_XII_PAGES_055-091 CH 03.indd 68 3/8/2020 2:44:19 PM