Table 6.4.2
Intermittent Fevers

	Malaria	Dengue
Causative Organism	Genus Plasmodium P. falciparum P. vivax P. ovale P. malarae	Genus Flavivirus DEN 1 DEN 2 DEN 3 DEN 4
Mode of transmission	Mosquito-borne protozoan disease Anopheles mosquito	Mosquito- borne viral disease Aedes aegypti and Aedes albopictus

6.4.2. Intermittent Fever

Intermittent fever usually present for some hours of the day and the rest of the time, temperature is normal. The increase in body temperature can occur same time each day or every few days in a repetitive pattern. Examples are Malaria, Dengue and Chikungunya.

A) Malaria

Malaria is a protozoan disease transmitted by the bite of the infected anopheles mosquito. Four species of genus ***Plasmodium*** causes malarial infection in humans.

Malaria comprises three stages

a) Cold stage (fever with chills)

b) Hot stage (temperature rises to >105°F and persists for 4 to 6 hours)

c) Sweating stage (fever subsides with sweating).


The clinical features are mild to severe fever, head ache, fatigue and muscle pain. The complications of malaria may be coma, convulsions, hyperpyrexia, hypoglycemia, liver damage, anemia and acute renal failure.

B) Dengue

Dengue is called as break bone fever. Dengue is caused by dengue virus which belongs to genus Flavivirus. It is spread through bite of infected Aedes mosquitoes, primarily Aedes aegypti and Aedes albopictus. It is characterized by nausea, vomiting, abdominal pain, rash, myalgia and bone pain and leukopenia. Two severe syndromes can be caused in young children. They are: Dengue Haemorrhagic Fever (DHF) and Dengue Shock Syndrome (DSS).



DHF, now called as "Severe Dengue" was recognized in 1950s in Philippines and Thailand.



Dengue fever symptoms and signs



Activity : 5

Hide and Seek!

Identify the areas in your house which attracts mosquitoes.

UNIT - 6 DIET IN FEVERS.indd 101 25-02-2019 15:35:35