diseased, they may rupture or there may be inadequate blood supply to brain resulting in stroke. This may be due to hypertension or fatty deposits in blood vessels of obese.

Type II Diabetes: It is an important contributor to morbidity and mortality in obese people. It is associated with insulin resistance and hyperinsulinaemia. A balanced diet, physical activity and drugs can control blood sugars.

Gall bladder disease: Obesity is one of the risk factors for formation of gallstones. The excess adipose tissue contain large amount of cholesterol. The super saturation of bile with cholesterol in obese makes them prone to formation of gallstones.


Cancer: Risk of cancers of colon, rectum and prostate increases in obese men, whereas obese women likely to develop cancer of breast, ovary, endometrial and cervix.

Infertility: Obese women are reported to suffer more from menstrual disorder, infertility and PCOS (Polycystic Ovary Syndrome), all of which tend to improve on reduction of weight.

Sleep Disorder: Commonly known as sleep apnoea is one of the common problems in obese men and women. Obesity causes narrowing of the upper airway when the person is in supine position.

Back pain and Arthritis: Abdominal obesity increases risk of back pain because of extra load on spinal column. The extra stress on weight bearing joints contributes to the development of osteoarthritis.

Psychological Problems: Obese people may be exposed to ridicule and discrimination in areas like employment, promotions and social interactions. This results in low self-esteem and depression leading to over eating for consolations.



(NIDDM-Non Insulin Dependent diabetes Mellitus)

Management of Obesity: Comprises the Following Three Approaches.

1. Dietary modifications
2. Physical activity
3. Lifestyle modifications

7.1.4. Dietary Principles:

A low calorie, restricted carbohydrate, high protein, restricted fat, normal vitamins and minerals (except sodium), liberal fluid and high fibre diet is suggested.

Energy: A low calorie diet helps the body to go into negative energy balance. This promotes use of stored fat as opposed to calories from food. 20 Kcal/kg ideal body weight for sedentary lifestyle and 25 Kcal / kg ideal body weight for moderate lifestyle is prescribed.

Carbohydrates: Milled and refined grains and foods made with refined grains like white rice, white bread, white pasta, processed breakfast cereals, potatoes and sugary drinks are rich in rapidly digested carbohydrate having high glycemic index and glycemic load. These kinds of carbohydrate increase the blood glucose and insulin levels rapidly, hence it causes hunger spikes leading to overeating and resulting in weight gain and consumption should be restricted.

UNIT - 7 OBESITY AND UNDERWEIGHT.indd 114 25-02-2019 15:36:09