Diagnosis	A1C (percent)	Fasting plasma glucose (FPG)[a]	Oral glucose tolerance test (OGTT)[b]	Random plasma glucose test (RPG)[a]
Normal	below 5.7	99 or below	139 or below	
Prediabetes	5.7 to 6.4	100 to 125	140 to 199	
Diabetes	6.5 or above	126 or above	200 or above	200 or above

(a) Glucose values are in milligrams per deci liter, or mg/dL.

(b) At 2 hours after drinking 75 grams of glucose.

Source: Adapted from American Diabetes Association. Classification and diagnosis of diabetes. Diabetes Care.2016;39(1):S14–S20, tables 2.1, 2.3.

Both methods are dependent upon a normal glomerular filtration rate (120ml/min) and renal threshold for glucose reabsorption by the tubules (180mg/dl).

Glycosylated Hemoglobin or 'Hemoglobin A1c' Test:

The Hemoglobin A1c test measures how high the person's blood sugar levels have been over the last 120 days. The test involves drawing a blood sample, and is the best way to measure blood sugar control in persons with diabetes. If the test results show 7% or less, the person has good blood glucose control. If the person has 8% or higher test results, their blood sugar has been too high for too long.



Activity : 3

Write the normal blood sugar levels for an adult ------ mg/dl.

9.7. Management

Although it is not possible to cure diabetes completely, diabetics can lead almost a normal life if they follow certain dos and don'ts scrupulously. Cooperation of the patient is very important in the management of Diabetes.

The main modes of treatment in Diabetes are:

1. Diet
2. Insulin
3. Drugs
4. Exercise
5. Education

9.7.1. Dietary management

A well designed meal plan is an important cornerstone in the management of diabetes mellitus.

Goals of nutritional therapy

- Achieve physiologic blood glucose levels.
- Maintain desirable plasma lipid levels.
- Reduce complications of diabetes mellitus.
- Attain and maintain desirable body weight.
- Meet energy needs in a timely manner.
- Individualize to preferences and food available.

The nutrition plan

Calories

Calorie requirement depends on the weight of a person. The calorie content of all diabetics should be set at a level which will permit them to maintain their desired body weight and in children and adolescents allow for a normal rate of growth and development.

UNIT 9_new.indd 148 25-02-2019 15:37:14