


Fig 10.2 Sodium rich foods

Potassium: Potassium is found naturally in almost all foods particularly in fruits, vegetables and meats. Potassium content can be reduced in vegetables by cooking in excess water and then discarding the water. This is called leeching. Spices and condiments can be used in small quantities as they are rich in potassium. The following foods which are high in potassium are restricted in a patient suffering from glomerulonephritis:




Fig 10.3 Potassium rich foods

- Nuts
- Jaggery
- Instant coffee
- Chocolate
- Cocoa powder
- Amaranth
- Spinach
- Colocasia
- Yam
- Mango
- Sweet lime
- Amla
- Lemon

Phosphorus: Eating foods high in phosphorus will raise the phosphorus in blood and this can cause calcium to be pulled from bones. This will make bones weak and cause them to break easily. The following foods which are rich in phosphorus should be restricted in the diet:




Fig 10.4 Phosphorus rich foods

- Milk
- Cheese
- Nuts
- Bakery products
- Pulses
- Mango
- Amla
- Lemon
- Eggs

Foods permitted and avoided in glomerulonephritis

Foods permitted

- Rice
- Milk or milk products, not more than 300-400 ml

Unit 10 Diet in kidney diseases Final 12-12-2018.indd 164 25-02-2019 15:37:56