- Vegetables, cooked in large amount of water with excess of water that is drained out
- Potato, sweet potato or yam
- Sugar
- Sago
- Gourd vegetables

Foods avoided

- Pulses
- Soups
- Meat,
- Fish
- Fruits rich in potassium
- Dried fruits
- Nuts
- Condiments and spices
- Papads and pickles



Activity : 2

Justify why the following foods are avoided or restricted in glomerulonephritis.

- Meat
- Milk
- Pulses
- Baked foods
- Nuts
- Jaggery
- Pickles
- Dried fruits
- Chocolate
- Eggs



Activity : 3

List any 5 foods that can be included in a diet for a patient suffering from glomerulonephritis.

1.

2.

3.

4.

5.

10.4. KIDNEY STONES (UROLITHIASIS)

Renal stones are prevalent between 35-40 years of age with the incidence declining after the age of 50. Urolithiasis or renal calculi can be found lodged in the urinary tract namely, kidney, ureters, bladder or urethra. The essential phase in stone formation is condensation of specific molecules within the collecting system. These molecules of mucopolysaccharides and mucoproteins form a matrix united by a strong chemical bond, with deposition of crystals in them as a secondary phenomenon.

10.4.1.Causes

The occurrence of kidney stones may be an outcome of different nutritional status, dietary habits and environmental factors such as temperature and humidity. The causes of renal stones are:

- **Heredity:** Some of the blood relations of patients with kidney stones develop the same problem.
- **Climate:** Those residing in hot tropical climates are prone to develop urinary calculi. During hot weather, less urine is formed as more water is lost through perspiration.
- **Fluids:** In the tropics, a lot of fluid is lost through perspiration, and so crystals are easily precipitated in the concentrated urine.
- **Vitamin B complex:** The metabolites excreted in the urine are significantly altered by the vitamin content in the diet. The administration of tryptophan increases, while vitamin B6 decreases urinary oxalate excretion.
- **Vitamin D and Calcium:** Excessive urinary excretion of calcium predisposes to formation of calcium stones. Vitamin D increases calcium absorption from the intestinal tract.

Unit 10 Diet in kidney diseases Final 12-12-2018.indd 165 25-02-2019 15:37:57