Table 11.1 : Sodium content of foods per 100 grams

<25 mg Low	25–50 mg Moderate	50–100 mg Moderately High	>100 mg High
Amla	Raisins	Cauliflower	Amaranth
Bitter gourd	Broad beans	Field beans	Bacon
Bottle gourd	Carrots	Fenugreek	Egg
Brinjal	Reddish (white)	Lettuce	Lobster
Cabbage	Black gram dal	Beetroot	
Lady's finger	Green gram dal	Water melon	
Colocasia	Red gram dal	Bengal gram dal	
Cucumber	Lentil	Red gram tender	
French beans	whole Bengal gram	Liver	
Peas	whole Banana	Prawns	
Onion	Pineapple	Beef	
Potato	Apple	Chicken	
Tomato	Mutton		
Yam			
Cow pea			
Horse gram			
Ragi			
Vermicelli			
Semolina			
Wheat			
Maida			
Milk			
Grapes			
Sweetlime			
Papaya			
Orange			
Sapota			

Source: Indian guidelines on hypertension-III – 2013. JAPI. 2013 Feb;61(2) suppl:16–19

sources of protein are high in saturated fat and therefore should be included in limited quantities.

- **Carbohydrate:** Simple Carbohydrates like sugar should be avoided. Complex carbohydrate from whole grains, pulses,

UNIT - 11.indd 179 25-02-2019 15:38:19