
Jejunostomy



Infusion techniques :

- **Continous method** - It implies continuous feeding through tubes by using a gravity flow or infusion pump.
- **Cyclic method** - This implies following a cycle of infusion by " off and on " periods. For example infusion is on for 8 hours and then is stopped for 16 hours.
- **Interrupted method** – this is like cyclic method but with regular interruptions like 6 hours on and 6 hours off periods.
- **Bolus method** – when a large feed is administered in a short time for example 250 cc in 15 minutes.

Types of enteral feeds :

1. **Natural liquid feeds**

   These comprise milk – (whole or skimmed), egg mixtures, sugar, molasses, strained cooked cereals, strained fruit juices, coconut water, vegetable oil, cream, non fat dry milk.

2. **Blenderized feeds**

   Foods which cannot be swallowed are blended in a high speed blender into a smooth paste which is reconstituted with water to make a thin liquid.

3. **Commercial polymeric mixtures**

   These contain all the carbohydrate, protein, fat, water, electrolytes, micro nutrients and fibre required by a stable patient

4. **Elemental feeds**

   They are commercially predigested mixtures of amino acids , dextrins , sugars, electrolytes , vitamins and minerals with small amount of fat .









UNIT - 5 Therapeutic diets.indd 89 25-02-2019 15:34:59