2 **Genetic factors:** It is common in persons with blood group 'O'.


3) **Gender:** Men are affected 2-3 times more frequently than woman.

4) **Age:** The incidence is high between 20 and 40 years

5) **Stress:** People who are highly nervous and emotional are more susceptible.

6) **Potential irritant substance:** Caffeine, chillies, pepper, ginger, meat soup and strong tea or coffee and protein rich foods increase the secretion of hydrochloric acid

7) **Emergency injuries :**Stress ulcers occur in conjunction with emergency injuries such as burns.



Symptoms of Ulcer

- Epigastric pain in the upper abdomen.
- Hemorrhage-an escape of blood from a ruptured blood vessel.
- Hypermotility- excessive motility of the gastrointestinal tract.
- Painful hunger contractions.
- Flatulence.
- Weight loss
- Iron deficiency.

Principles of diet

High energy, high protein, moderate fat and low fibre

Regular balanced diet should be given to support the primary medical management of

1) tissue healing and

2) maintenance of structural and functional integrity of tissues.

Dietary Management

Energy: Increased energy intake is necessary. If the patient is at bed rest, energy is not required for activity but needed to make-up the extra demands.

Protein: A high protein diet promotes healing. Milk and milk products should be included as a source of nutrient factor for healing process. Eggs and other high protein foods can be included.

Fat: Moderate amounts of fat to help to suppress gastric secretions and motility.

Carbohydrates: They are used to meet the energy needs. Foods containing high fibre and vegetables which are high in insoluble fibre are restricted.

Fibre: Soluble fibre can be given. Certain raw fruits and vegetables high in fibre are restricted.

Gastritis is the inflammation and corrosion of the inner lining of the stomach called mucosa as a result of an immune response to infection or external injury.

XII Std - Home Science EM_Unit 1.indd 9 20-04-2020 6.42.48 PM