Minerals: Restriction of sodium as common salt is helpful in weight reducing diet as excess sodium leads to retention of fluid.

Fluid: Fluid can be taken liberally. Also a glass of water before meal helps to cut down food intake.

High Fibre: High fibre low calorie foods like green leafy vegetables, fruits, vegetables salads, whole grain cereals and pulses can be included in the diet.

Advantages of High Fibre Food

1. Low in calorie density
2. Foods like greens provide many vitamins and minerals (which are difficult to meet with restricted foods)
3. Give satiety
4. Help in regulating bowel movements
5. Reduce blood cholesterol
6. Promote chewing and decrease rate of ingestion.

A paleo diet typically includes lean meats, fish, fruits, vegetables, nuts and seeds.



The colour Orange is named after the Orange fruit, but before that, it was called geoluread (yellow-red)

Physical Exercise

A low calorie diet accompanied by moderate exercise will be effective in causing weight loss.

- Aerobic exercise directly increases the daily energy expenditure.
- Exercise will also preserve lean body mass
- It helps to regulate appetite and increases the basal metabolic rate
- Reduce stress related eating.

Complications of Obesity

- ***Physical disability***: Feet have to carry extra load complications like flat feet, osteo-arthritis of knee, hip and lumbar spine.
- ***Metabolic disorder***- (a) Diabetes. (b) Plasma cholesterol level. (c) Atherosclerosis. (d) Gout.
- ***Cardiovascular disorders***-: High blood pressure and varicose veins.
- ***Sleep apnea***.
- ***Prone to accidents***-: Falling down on slippery floor and crossing streets.
- ***Gastrointestinal disturbance***: Hiatal hernias and gall bladder disease.
- ***Osteoarthritis***.
- ***Cancer***: Cancers (colon, rectum and prostrate in men, uterus, biliary tract, breast and ovary in women), digestive tract diseases (gallstones, reflux, esophagitis) and skin disorders.

1.5.2 Underweight

According to the WHO classification of Body Mass Index (BMI) people whose BMI is less than 18.5 are considered as underweight.

Causes of Underweight

Starvation: This occurs due to famine conditions or an inadequate diet lacking in proteins or an attempt at reducing weight.

XII Std - Home Science EM_Unit 1.indd 18 20-04-2020 6.42.49 PM