Clinical symptoms

- Oedema
- Ascites (accumulation of fluid in the abdominal cavity)
- Proteinuria
- Distended abdomen due to fluid accumulation.
- Albuminuria (increased loss of urinary albumin)
- Malnourishment due to large protein loss that leads to tissues protein breakdown.

Principles of diet

High energy, moderate protein, salt and fluid restricted diet is advised.

Dietary management

Energy

A high intake of 30-40 g/kg ideal body weight is provided for tissue rebuilding. Food to be made more appetizing and soft in consistency to improve acceptability.

Protein

A protein intake of 0.8- 1.0 g/kg body weight/day given to meet the nutritional needs and prevent tissue protein breakdown. High quality proteins present in animal foods are preferred.

Sodium

Sodium is reduced to approximately 1 g/day to prevent oedema and increased to 3 g/day as oedema subsides.

Other minerals and vitamins

Iron and vitamin supplement may be helpful, while potassium restriction is not necessary. Potassium supplements are useful to replace losses with diuretic usage.

DO YOU KNOW?


Kidney transplantation is needed when the kidney function reduces to less than 10%. Donor kidneys are taken from either brain dead organ donors or family member of the recipients.

DO YOU KNOW?

Human Organ Transplantation Act (1994) in India allows only live transplants from close relative/voluntary donors (exceptional) to avoid organ trafficking.

1.8.3 Kidney Stones/ Urinary Calculi

Kidney stones, also called urinary calculi /nephrolithiasis may be found in the kidney, ureter, bladder or urethra.



The basic cause of kidney stones is unknown, but many factors relating to the nature of the urine itself or conditions of the urinary tract environment contribute to their formation.

XII Std - Home Science EM_Unit 1.indd 38 20-04-2020 6.42.54 PM