3.4.1 Common Adulterants in food and their Harmful effects

Sl. No.	Name of the food article	Adulterant	Simple method to detect the adulterant	Harmful Effects
1.	Ghee or butter	Vanaspathi	Take one teaspoonful of ghee or butter with equal quantity of concentrated hydrochloric acid in atest tube and add a pinch of cane sugar to it . Shake well for one minute and test it after 5 minutes. Appearance of crimson color shows the presence of vanaspathi	____
2.	Milk	Water	Lactometer reading should not be less than 1.026	Stomach disorder
			Drop of pure milk flows slowly leaving a white trail behind whereas adulterated milk with water will flow immediately.	___
3.	Khoa	Starch	Add tincture of iodine. Blue color shows the presence of starch.	Less - nutritive value
4.	Dhals	Kesari dhal	Add 50 ml of dilute hydrochloric acid to the dhal and simmer it for 15 minutes. Pink color shows the presence of kesari dhal.	Stomach pain, ulcer
5.	Hing	Soap stone or other earthy matter	Shake with water , soap stone or earthy matter will settle down.	Dysentry
6.	Tea leaves	Exhausted tea or black or Bengal gram dhal husk with color	Tea leaves sprinkled on wet filter paper would immediately release added color.	Liver Disorder
7.	Sugar	Chalk Powder	Dissolve in a glass of water, chalk will settle down at the bottom	Stomach – Disorder
8.	Chilli powder	Stones Brick Powder, Sawdust	Any grittiness present may be felt by tapping. The sediment at the bottom of glass confirms the presence of brick powder or sand.	Stomach problems

XII Std - Home Science EM_Unit 3.indd 83 23-03-2020 3.29.40 PM