causing food poisoning. S. enteritids, S. typhimurium, S. Thompson are the most frequent but other types have also been isolated in outbreaks of food poisoning. Salmonella food poisoning causes gastro enteritis since only the alimentary tract is involved and there is no general invasion of the body.

Signs and symptoms: The condition is characterized by nausea, vomiting, abdominal discomfort, headache and later diarrhoea. The symptoms appear 12-26 hours after the ingestion of the contaminated food. In most cases these symptoms subside in a few days and the patients recover completely in 6-8 days. Mortality rate is very low.

Foods involved: The common foods that give rise to this type of food poisoning are meat, milk, fish or eggs, the contamination being from the animal itself or from external sources. Eggs from infected ducks have been the cause of several outbreaks, many types of Salmonella have been isolated from dried egg powder and egg albumen.


Staphylococcus Food Poisoning:

Causative agents: It is due to Staphylococcus aureus and strains of this organism produce an exotoxin which is responsible for a number of outbreaks of food poisoning characterized by vomiting and diarrhea. The sources include human or animal origin. The nasal passages of many persons contain a large number of these organisms, boils and infected wounds may also be sources of contaminating food.

Signs and Symptoms: The duration of the condition caused by staphylococcus food poisoning is only for a day or two. Mortality is extremely low.

FOOD POISONING SYMPTOMS

B3MUWY



XII Std - Home Science EM_Unit 3.indd 86 23-03-2020 3.29.40 PM