The *SDR* can be calculated with respect to gender also. For example, *SDR* for males in a given region during a given period can be calculated as

$$SDR_{Male} = \frac{D_{Male}}{P_{Male}} \times 1000 .$$

Here, D_{Male} is the number of male deaths in the region during the specified period, and P_{Male} is the male population size in the region during the specified period.

(iii) Infant Mortality Rate

'Infant' means a baby of age less than a year. **Infant Mortality Rate** (*IMR*) is defined as the number of infant deaths *per thousand* live births in a period or children die before they attain age of one year. The following formula is used to calculate infant mortality rate

$$IMR = \frac{D_{Infant}}{P_{Infant}} \times 1000$$

where

D_{Infant} denotes the number of deaths of infants in a population during a period, and

P_{Infant} denotes the number of live births in the population during the period.

Child Mortality Rate: Child mortality is the death of a child before the child's fifth birth day, measured as the under 5 child mortality rate (*U5MR*).

Example 8.5

The number of live births recorded and the number of infants died in a town during a given period are respectively 400 and 25. Calculate, from these information, the infant mortality rate of the town for the period.

Solution:

The *IMR* of the town is given by

$$IMR = \frac{25}{400} \times 1000$$

$IMR = 62.50$ *per thousand.*

8.1.4 Life Table and Its Applications

A **Life Table** is a presentation of summary of the mortality experiences of a community during a given period in the form of a table. The Life Table exhibits the number of persons living and dying at each age, on the basis of the experience of a ***cohort***. It also gives the probability of dying and living separately. The Life Table tells the life history of a ***cohort***.

Cohort is a group of individuals who born at the same time and experienced the same mortality conditions.

12th_Statistics_EM_Unit_8.indd 216 1/28/2020 6:38:10 PM