Deficiency

Vitamin D deficiency affects the mineralization of bones and teeth.

-Rickets, osteomalacia

Vitamin E (Tocopherol)

Vitamin E is an essential, fat – soluble vitamin that includes eight naturally occurring compounds in two classes designated as tocopherols and tocotrienols.

Vitamin E is an effective chain – breaking, lipid – soluble antioxidant in biological membranes, and aids in membrane stability.



Sources of Vitamin E

Function

- Vitamin E is a powerful antioxidant.
- Antioxidant capability helps to prevent degenerative diseases.
- Vitamin E is also useful in preventing blood clots forming and promotes fertility.
- An increase in stamina and endurance is also attributed to vitamin E.
- Vitamin E is also used to great effect for skin treatments.

Deficiency of vitamin E


Deficiency of vitamin E is a not common, and the symptoms not very clear cut, but may include fatigue, inflamed varicose veins, slow wound healing, premature ageing and sub – fertility.

Vitamin K

Vitamin K can be produced in the intestines and this function is improved with the presence of cultured milk, like yogurt, in the diet.

Functions

- The major function of vitamin K is to promote coagulation of blood after injury, thereby preventing haemorrhage.
- Vitamin K is necessary for the synthesis of prothrombin, an inactive form of thrombin.
- It is involved in the formation of prothrombin.
- It is also involved in bone formation and repair.



Food Source



Sources of Vitamin K

The best dietary sources of this vitamin are green leafy vegetables, cheese and liver.

Deficiency

Increased tendency to haemorrhage defective blood clotting

Unit_5_Applied_nutrition_Gen_Nur.indd 121 04-03-2020 17:12:19