and glycosuria accompanied by polyuria, polydipsia, polyphagia, and if untreated for a long time by loss of weight.

Hypertension

Hypertension develops due to some reason, there is increased resistance to the normal circulation of blood. This may be temporary, as the result of exercise, pregnancy, or menopause; it may be permanent, associated with atherosclerosis, cardiac failure or chronic kidney disease.

Notes

1. Salt should not be added in cooking
2. One multivitamin tablet providing the daily requirements of all essential vitamins should be included along with the diet
3. Strong coffee is not permitted.

Table 14: Daily menu for Hypertension

Food stuffs	Vegetarian	Non – vegetarian
Milled cereals	200	200
Skin mil (fluid)	1000	1000
Pulses	60	40
Cheese	60	20
Meat or fish	-	20
Leafy vegetables	100	100
Other vegetables	100	100
Fruits	300	300
Vegetable oils (sesame or safflower or sunflower seed oil)	60	60
Sugar and jam	40	40

Coronary Heart Diseases

A majority of coronary heart diseases are due to coronary antheroma (A fatty deposit in the inner lining of an artery) and atherosclerosis, are used to denote conditions in which lipid is deposited in the intima of the blood vessels.

Calories – 1800 – 2000 Kcal; Fat (rich in essential fatty acids) – 60 gm; Proteins – 50 – 60 gm

Note: one multivitamin tablet providing the daily requirements of all essential vitamins should be given daily.

Table 15: Daily menu for coronary heart disease

Vegetarian	Non – vegetarian
Morning	
Weak tea	Weak tea
Breakfast	
Corn flakes with skimme milk or bread of idly – 1 serving Cheese – 2 slices Fruits – 1 serving Weak tea – 1 cup	Corn flakes with skimmed milk or bread or idly – 1 serving Cheese – 2 slice Fruits – 1 serving Weak tea – 1 cup
Lunch	
Cooked rice or chapatti – 1 serving Dal soup – 1 cup Cooked vegetables – 1 serving Cheese – 2 slices Curds – 2 cups Fruits – 1 serving Skimmed milk pudding – half cup	Cooked rice or chapatti – 1 serving Mutton soup – 1 cup Mutton or fish curry – 1 serving Curds – 1 cup Fruits – 1 serving Skimmed milk pudding – half cup
Tea	
Biscuits – 2 Fruits – 1 serving Weak tea – 1 cup	Biscuits – 2 Fruits – 1 serving Weak tea – 1 cup
Dinner	
Similar to Lunch	

Chronic Renal Failure

Chronic Renal Failure occurs in glomerulonephritis, and chronic pyelonephritis. In this condition, the glomeruli and tubules are heavily damaged.

Unit_5_Applied_nutrition_Gen_Nur.indd 134 04-03-2020 17:12:20