Diet

The diet in chronic uratemia (presence of abnormal of uric acid salts in the blood) should be below in protein depending on the urea content of blood. The pattern of diet and the daily menu for an adult suffering from chronic renal failure are given in Tables 16 and 17.

Table: 16 Diet for an adult suffering from chronic renal failure

(Calories 1500 – 2000 Kcal; Proteins 15 – 20 gm; Fat 40 – 50 gm, Carbohydrates 300 – 40 gm; Salt free)

Foodstuffs	gm/caput/day
Cereals	200
Vegetables (potato, carrot)	100
Milk	200
Egg	30
Butter and vegetable fats	40 – 50
Sugar	50
Glucose	200
Fruit juice	1000 ml

Table 17: Daily menu for chronic renal failure

Morning	Fruit juice with glucose – 1 glass
Breakfast	Bread – 2 slice Butter – 2 teaspoons Jam – 2 teaspoons Fruit juice with glucose – 1 glass
Lunch	Cooked rice or bread – 1 serving Butter or ghee – 2 teaspoons Potato and carrot curry – 1 serving Curds – 1cup Fruit juice with glucose – 1 glass
Tea	Biscuits – 3 Butter – 1 teaspoon Fruit juice with glucose – 1 glass
Dinner	Similar to Lunch

Note: Common salt should not be added during or after cooking. One multivitamin tablet providing the daily requirements should be taken

Peptic Ulcer

Chronic ulcer formed in the region of the gastrointestinal tract where the gastric juice comes in direct contact with the mucous membrane is known as peptic ulcer. Such ulcers usually occur in the duodenum (duodenal ulcer) and in the stomach (gastric ulcer). Duodenal ulcer is associated with hyperacidity while gastric ulcer is not.

Diet in Peptic Ulcer

The diet is the most important factor in the treatment of peptic ulcer. Three main types of diets have been used in the treatment of peptic ulcer.

Table 18: Types of therapeutic diet for peptic ulcer

Name of Diet	Ingredients
Sippy's diet and its modification by Hurst	Hourly feeds of milk, cream and olive oil with antacid medication
Lenhartz diet	Fluid diet based on milk and eggs
Meulengracht diet	A mixed bland diet containing milk, egg, meat and fish (Minced and strained) given once in 2 hours.

Sippy's diet consisting of milk and olive oil was used widely. Since it is monotonous and not adequate to meet the calorie needs, it has been replaced by a bland diet containing large quantities of milk.

Table 19: Diet for adult suffering from peptic ulcer (gm/caput/day)

Foodstuffs	Vegetarian	Non – vegetarian
Milled rice and refined wheat flour, or bread and biscuit	200	200

Unit_5_Applied_nutrition_Gen_Nur.indd 135 04-03-2020 17:12:20