Dal (decuticled split legumes)	50	50
Milk	1800	1500
Cheese	50	-
Eggs	-	30
Meat or lean fish	-	60
Fats and oils (2/3 as vegetable oils rich in essential fatty acids)	60	60
Sugar	60	60
Tender vegetables	50	50
Potato	50	50
Fleshy fruits (banana, apple and mango)	100	100

Note: One vitamin C tablet (100mg) may be taken once daily.

Table 20

Vegetarian	Non – vegetarian
Morning 6 am	
Milk – 2 cups (with 2 teaspoon sugar)	Milk – 2 cups (with 2 teaspoon sugar)
Breakfast 8 am	
Breads – 2 slices Butter – 2 teaspoons Cheese – 2 slices Milk – 1 cup	Breads – 2 slices Butter – 2 teaspoons Boiled egg – one Milk – 1 cup
10 am	
Milk – 2 cups (with sugar)	Milk – 2 cups (with sugar)
12 noon (Lunch)	
Cooked rice or bread – 1 serving Smashed dal – 1 cup Cheese – 2 slices Boiled potato – 2 Milk pudding – 1 cup	Cooked rice or bread – 1 serving Smashed dal – 1 cup Cooked minced meat – 1 serving Boiled potato – two Milk pudding – 1 cup
2 pm	
Milk – 1 cup	Milk – 1 cup
4 pm	
Biscuits – 2 Milk (with sugar) – 1 cup	Biscuits – 2 Milk (with sugar) – 1 cup
6 pm	
Milk (with sugar) – 2 cups	Milk (with sugar) – 2 cups
8 pm (Dinner)	
Same as Lunch	Same as Lunch
10 pm	
Milk – 1 cup	Milk – 1 cup

Table 21: Foods permitted and not permitted in peptic ulcer

Permitted	Not permitted
1. Milled cereals (milled rice, refined wheat flour and semolina) 2. Dal (decuticled split legumes) 3. Milk 4. Eggs 5. Tender vegetables free from fiber 6. Meat and fish in limited amounts 7. Fruits without fiber 8. Fats and oils 9. Sugar	1. Whole cereals and whole millets 2. Legumes with husk (Bengal gram, field bean, cow gram, dried pea, etc.). 3. Vegetables containing fiber, raw onions, garlic, etc. 4. Spices and condiments and pickles 5. Meat soups and extractives 6. Fried foods 7. Alcohol and alcoholic drinks 8. Chewing betel leaves, arecanut and tobacco

5.11 Adulteration of Food Additives

Food additives are defined as non-nutritious substances which are added intentionally to food generally in small

Unit_5_Applied_nutrition_Gen_Nur.indd 136 04-03-2020 17:12:20