DO YOU KNOW?

These are some of the beneficial pregnancy tips: -

- Eat 5 to 6 meal a day
- Take a prenatal vitamin each day as directed by the doctor
- Drink plenty of fluids
- Take adequate rest and sleep
- Do proper exercise
- Wear comfortable clothes
- Don't take over the counter drugs
- Wear safety belts while traveling
- Avoid exposure to smoking

7.4 High Risk Pregnancy

Definition

High Risk Pregnancy is a pregnancy complicated by a disease or a disorder that may endanger the life or affect the health of the mother, the fetus or the newborn

Maternal Death Incidence

A maternal death is death of a Woman while pregnant or within 42 days of termination of pregnancy irrespective of the duration and the site of the pregnancy, from any course related to or aggravated by the pregnancy or in management but not from accidental or incidental causes. - WHO

- 20-25% deaths occur during pregnancy.
- 40-50% deaths occur during labour and delivery.
- 25-40% deaths occur after childbirth (More during the first seven days)
- Annually, 585,000 women die of pregnancy related complications

High risk mothers are

- Women below 18 years, over 35 years in primigravida
- Women who had four or more pregnancies
- Short Structure (height <145 cm and below)
- Twins
- Aneamia
- Previous Abortion, Intra uterine death.
- Malnutrition mother.

Placenta Praevia

DEFINITION: Placenta Praevia is a condition where the placenta is implanted completely or partially in the lower part of the uterus.

Cause: Unknown


Risk factors: Multiparity, multiple gestations, previous uterine surgery

Manifestations: Painless, bright red bleeding > 20th week; episodic, starts without warning, stops & starts again.

Vaginal examination is contra - indicated.

Types of Placenta Praevia

- First degree (Type I): Low lying Placenta the lower edge of the placenta reaches the lower uterine segment but not the internal cervical os.
- Second degree (Type II): Marginal the lower edge of the placenta reaches the margin of the internal os but does not cover it.
- Third degree (Type III): Incomplete or partial the placenta covers the internal os partially.
- Fourth degree (Type IV): Total placenta covers the internal os completely.



Unit_7_Midwifery_Gen_Nur.indd 157 09-03-2020 14:51:12