

Management

Management depends upon gestational age, amount of bleeding and fetal condition.

- Monitor Fetal Heart Rate, maternal Vital signs
- Intra Venus Fluid administration
- O_2 administration
- Assess intake and output, amount of bleeding
- Do complete Blood count and Rh factor test (CBC), Type and cross match for transfusion.
- Ultrasound
- No pelvic exams
- No vaginal delivery- may lead to haemorrhage
- Prepare for caesarean section

Prognosis: depends on amount of bleeding & gestational age

Abruptio Placenta

Abruptio Placenta: Premature separation of normally situated placenta.


Cause: Unknown

Types of Abruptio placenta:

1. **Concealed** - The blood collects behind the separated placenta or collected in between the membranes and decidua. (Blood is not visible outside) Rare type.
2. **Revealed** - Following separation of the placenta, the blood comes out of the cervical canal to be visible externally (commenest type)

Risk factors

- Smoking
- Short umbilical cord
- Advanced maternal age
- HTN
- PIH
- Cocaine use
- Trauma to or near abdomen.



Manifestations: Tenderness mild to severe constant pain; mild to moderate bleeding depending on degree of separation.

Management

- Monitor Fetal Heart Rate, maternal Vital signs
- Intra Venous Fluid administration
- O_2 administration
- Assess intake and output, amount of bleeding
- Do complete Blood count and RH factor test (CBC), Type and cross match for transfusion.
- Ultrasound
- No pelvic exams
- No vaginal delivery- may lead to haemorrhage
- Prepare for caesarean section

Unit_7_Midwifery_Gen_Nur.indd 158 09-03-2020 14:51:12