Symptoms

- Colicky, cramping pain in lower abdomen on affected side
- Tubal rupture: sudden/sharp/steady pain before diffusing throughout pelvic region
- Heavy bleeding causes shoulder pain, rectal pressure
- Dizziness/weakness - If tube ruptures, weak pulse, clammy skin, fainting. Assess for s/s shock.

Diagnosis

Estimation of Beta hCG (more than 1500 IU/L

- Ultrasonogram

Treatment

- Immediate surgery to remove/repair tube.
- If no rupture, Methotrexate - stops cellular division in fetus; causes cell death. Conceptus expelled with bleeding.

Hypertension In Pregnancy

Global cause of maternal/fetal morbidity & mortality. Responsible for 76,000 deaths/ year. Normotensive patient may become hypertensive in late pregnancy, during labor, or 24 hours postpartum.

Pre-Eclampsia

Defined As

Pre-Eclamsia is characterised by hyper tention protrinuria and oedema.

Dangers of Pregnancy Induced Hyper tention (PIH)

- BP ≥ 140/90 mmHg
- Systolic ↑ of 30mm Hg > pre-pregnancy. levels
- Diastolic ↑ of 15mm Hg > pre pregnancy. levels.
- Presents with HTN (Hypertension), proteinuria, edema of face, hands, ankles.
- Can occur anytime > 20th week of pregnancy.
- Usually occurs closer to due date. Will not resolve until birth.

General Signs of PRE-ECLAMPSIA

- Rapid weight gain; swelling of arms/face
- Headache; vision changes (blurred vision, seeing double, seeing spots)
- Dizziness/faintness/ringing in ears/ confusion; seizures
- Abdominal pain, ↓ production of urine; nausea, vomiting.
- Alarming signs:
  U - Urinary output dimished
  S - Sleep disturbance
  H - Headache
  E - Epigastric pain and eye symptoms.

Eclampsia

Seizures or coma due to hypertensive encephalopathy

Incidence

- Most serious complication.
- Affects ~ 0.2% pregnancy.
- Major cause of maternal death due to intracranial hemorrhage.
- Maternal mortality rate is 8-36%.

Risk factors

- < Age 20 years or > 40 years
- Twins, triplets
- Primigravida
- Molar pregnancy
- Preexisting HTN, Diabetes mellitus
- Renal or vascular disease
- Previous history of preeclampsia/eclampsia

Causes: Unknown.

Management: Usually only cure is termination of pregnancy. It depends upon symptoms.

Mild preeclampsia

- Bedrest
- Monitor at home or hospital.

Unit_7_Midwifery_Gen_Nur.indd 160 09-03-2020 14:51:13