Progestin releasing device thicken the cervical mucus and thin the endometrial lining thus prevents conception.

Types:

**Copper**

- Copper - T 380 A
- Nova T
- Multiload 375

**Silastic**

**Hormone releasing**

- Progestasert
- Levonova / Mirena

**Insertion:**

1. Bladder empty, lithotomy position.
2. Posterior vaginal speculum is introduced in vagina and cervix are cleansed by antiseptic lotion.
3. Anterior lip grasped by Allis forceps. Uterine sound passed through the cervical canal to normal position of uterus and length of cavity.
4. Insert the device through the cervical Os up to the fundus and after positioning, inserter is withdrawn keeping the plunger in position.
5. Excess of nylon thread is to be cut.

| Advantages | Disadvantages |
|---|---|
| • Highly effective in preventing pregnancy.<br>• Immediately effective after insertion.<br>• Long term protection is provided with copper T.<br>• Can be used for emergency post coital contraception.<br>• Do not affect breast feeding. | • Risk for pelvic inflammatory diseases.<br>• Dysmenorrhoea and menorrhea within first few months after insertion.<br>• IUD's offer no protection against STD's. |

### Side effects

- Cramping
- Abnormal menstrual bleeding

### Complications

- Uterine and pelvic infection
- Expulsion
- Uterine perforation
- Ectopic pregnancy

## IV. Hormonal Contraceptives

### Combined oral contraceptives

They consist of synthetic oestrogen and progestin preparations, act by suppressing lactation, thickening of cervical mucus and alteration of endometrium.

### Contraindication

| Absolute | Relative |
|---|---|
| • Breast carcinoma<br>• History of DVT, Pulmonary embolism<br>• Acute liver diseases<br>• Concurrent treatment with rifambicin<br>• Familial hyperlipidemia<br>• Pregnancy | • Smoking<br>• Age > 35yrs<br>• Obesity<br>• Hypertension<br>• Breast feeding |

### Side effects

- Amenorrhoea
- Breast fullness or tenderness
- Depression, severe vascular headache
- Hypertension
- Spoting or intermenstrual bleeding

### Missed pills

- If one pill is missed, instruct the patient to take two pills at the next scheduled time and complete the pack as usual.
- If two or more consecutive pills are missed, instruct the patient to finish the package of pills.



Unit_7_Midwifery_Gen_Nur.indd 170 09-03-2020 14:51:13