Progestron Only Pills

They cause suppression of ovulation, thickening of cervical mucus and alteration of the endometrium.

Advantages	Disadvantages
• Pill is rapidly effective • Fertility returns immediately when the pill is discontinued • Does not affect breast feeding • May improve anemia • Menstrual flow is decreased • Protects against endometrial cancer and pelvic inflammatory diseases	• Amenorrhoea • Intermenstrual bleeding • Must be taken everyday

Progestron only Injectable Contraceptives

They cause suppression of ovulation, thickening of cervical mucus, alteration of endometrium and change in tubal motility.

Advantages	Disadvantages
• Very & rapidly effective and long acting • Can be used by women more than 35 years.	• Return to fertility is delayed up to 5 – 7 months • Some patients experience weight gain, irregular bleeding, amenorrhea & excessive bleeding.

Implants (Norplant)

It consists of thin, flexible capsules filled with levonorgestrel that are inserted under the skin of women's arm. They cause suppression of ovulation, thickening of cervical mucosa, alteration of the endometrium and change in tubal motility.

Advantages	Disadvantages
• Highly effective and acts within 24 hours placement • Long term protection (3 – 5 yrs) • Immediate return of fertility on removal.	• Requires minor surgical procedure to insert and removal. • Provides no protection against STD's.

Advantages	Disadvantages
• Highly effective in preventing pregnancy. • Protection against endometrial cancer. • Benign breast disease, ectopic pregnancy and iron deficiency anaemia are also reduced.	• No protection against HBV, HIV or other STD's. • Must be taken every day.

Side effects

- Mastalgia
- Breast tenderness
- Weight gain or loss
- Irregular bleeding or spotting
- Amenorrhoea
- tHirsutism
- Hair loss

Permanent Method

Surgical Sterilization

It is appropriate to couples who desire permanent sterilization.

1. FEMALE: Tubal ligation OR Tubectomy

Types

- Puerperal sterilization
- 2. Interval tubal ligation:
- Minilaprotomy: The Pomeroy procedure performed using a 3 to 4 cm suprapubic incision under local anaesthesia with sedation.

Unit_7_Midwifery_Gen_Nur.indd 171 09-03-2020 14:51:13