Universal Immunization Programme

The **World Health Organization** officially launched a global immunization programme, known as **Expanded Programme on Immunization (EPI)** in May 1974, to protect all children of the world against six vaccine –preventable diseases. They are namely diphtheria, whooping cough, tetanus, polio, tuberculosis and measles. EPI was launched in India in January 1978. This programme is now called as **Universal Immunization Programme (UIP)** in India.

National Immunization Schedule for Infants and Children in India



BCG Vaccination

In addition to the above mentioned vaccines, Rota Virus vaccine & Japanese Encephalitis vaccine is given in selected states.

Diseases and its vaccine

S. N	Vaccine	Protection against
1	BCG	Tuberculosis
2	Hepatitis B	Hepatitis B, liver infection
3	Oral Polio Vaccine	Polio
4	Inactivated Polio Vaccine	
5	Pentavalent vaccine	Diptheria, Pertussis, Tetanus, Hepatitis B &Haemophilus Influenza B
6	Measles Vaccine	Measles
7	DPT Vaccine	Diptheria, Pertussis, Tetanus
8	TT (Tetanus Toxoid)	Tetanus
9	Rota virus Vaccine	Diarrhoea
10	Japanese Encephalitis Vaccine	Japanese Encephalitis - a brain infection

Vaccine	When to give	Dose	Route	Site
BCG (Bacilli galmetteGurein)	At birth	0.05ml	Intra dermal	Left Upper arm
Hepatitis B birth dose	At birth	0.5ml	Intra muscular	Left Antero lateral side of mid thigh
OPV zero dose	At birth	2 drops	Oral	
OPV 1, 2, &3 (Oral Polio Vaccine)	At 6, 10 & 14 weeks	2 drops	Oral	
IPV (Inactivated Polio Vaccine)	At 14 weeks	0.5ml	Intra muscular	Right-Antero lateral side of mid thigh
Pentavalent 1,2 & 3 (Diptheria, Pertussis, Tetanus, Hepatitis B &HIB)	At 6, 10 & 14 weeks	0.5ml	Intra muscular	Left Antero lateral side of mid thigh
Measles – 1st Dose	At 9 completed months	0.5 ml	Subcutaneous	Right Upper arm
DPT Booster-1	16-24 months	0.5ml	Intra muscular	Left Antero lateral side of mid thigh
Measles – 1st Booster Dose	16-24 months	0.5 ml	Subcutaneous	Right Upper arm
OPV Booster	16-24 months	2 drops	Oral	
DPT Booster-2	5-6 years	0.5ml	Intra muscular	Left Upper arm
Tetanus Toxoid (TT)	10 years & 16 years	0.5 ml	Intra muscular	Upper arm

Unit_8_Child_health_Gen_Nur.indd 184 09-03-2020 14:55:41