Mutism	Unable to speak	ஓசையற்ற/ஒலிநிறுத்தம்
Cognitive	Connected with thinking or conscious mental processes	அறிவாற்றல்
Relationship	State of being connected	உறவு
Intimate	Very close	நெருக்கமான
Therapeutic	Relating to the healing of disease	நோய்நீக்கநலம்சார்ந்த / குணப்படுத்தும் இயல்புடைய
Termination	Action of terminating something	முடிவிடம் / கடைமுடிவு
Media	The collective communication outlets	தகவல்தொடர்புசாதனம்
Anemia	A decrease in the total amount of red blood cells.	இரத்தசோகை
Retarded	Less advanced in physical, mental, or social development than usual age.	சரியில்லாத
Hinder	Make it difficult to do something	தடங்கல்செய் / இடையூறாக
Abuse	Treat with cruelty/violence. Use for a bad purpose	தவறானவகையில் பயன்படுத்துதல்
Disability	A physical or mental condition that limits a person's Activities	குறைப்பாடு / பலவீனம்
Head injury	Any trauma to the scalp, skull or brain	தலைக்காயம்
Illegible	Not clear enough to read	தெளிவில்லாத / விளங்காத / படிக்கமுடியாத
Genetic	Relating to genes or heredity	மரபணு
ADHD		அவதானக்குறை மிகையியக்க குறைபாடு.
Prevention	The action ofstopping something from happening	தடுத்தல் / நடைபெறாமல்செய்தல்
Rehabilitation	The action of restoring someone to health or normal life that has Been damaged.	மறுசீரமைப்பு / புனர்வாழ்வு

CLINICAL EXAMPLE

The Client with Severe Mental Retardation

Mr. X 18 years old, was born by a normal spontaneous vaginal delivery without any complications. Two months after birth, Mr. X developed a temperature of 105Â°F and had a grand mal seizure. He was admitted to the neonatal intensive care unit with the diagnosis of fever of undetermined origin (FUO). Diagnostic tests revealed the presence of encephalitis. MR. X recovered, but his parents were cautioned about the possibility of central nervous system damage because of the severity of his illness. During early childhood, ages 1 to 5, Mr. X was able to communicate with his parents to some extent but exhibited poor motor-skill development. He was unable to learn basic skills such as reading, writing, and arithmetic, and it became evident that he needed supervision in a controlled environment. Testing revealed Mr. X's IQ to be that of a person with severe retardation or mental impairment. At age 18, he continues to live at home with very supportive parents who have been able to teach him some self-care activities. He relates well to a pet cat, helps his mother with simple household chores, and helps his father with gardening and lawn care.

Unit_10_Mental_health_Gen_Nur.indd 242 09-03-2020 15:07:13