internationally. Health workers at all levels should be trained to identify and notify cases immediately to the local health authority.

3. **Early case finding** : An aggressive search for case (mild, moderate, severe) should be made in the community to be able to initiate prompt treatment.
4. **Establishment of treatment centers** : In the control of cholera, no time should be lost in providing treatment for the patients.
   1. Easily accessible treatment facilities in the community
   2. Treatment for dehydration
   - For Mild dehydration - treatment at home with oral rehydration fluid.
   - For Severely dehydrated patients requires intravenous fluids in a nearest treatment centre or hospital.
   - When it is endemic or threatening - mobile teams should be established.
3. Epidemiological investigations to study the extent of the outbreak and identify the modes of transmission.
4. Sanitation measures
   a. Water Sanitation
   b. Excreta Disposal - Health education messages should stress the proper use of such facilities.
   c. Food sanitation - Sale of foods under hygienic conditions. Health education regarding eating cooked hot food, and proper food handling techniques.
   d. Disinfection: Most effective disinfectant is bleaching powder. Clothes and personal items to be disinfected with disinfectant (dettol).
5. Health Education regarding Oral Rehydration therapy, Early diagnosis and adequate treatment and Food sanitation

Treatment

1. Replacement of fluids and Electrolytes.
2. Antibiotics
3. Anti diarrheal drugs
4. Rehydration therapy : The oral rehydration therapy and intra venous fluids (IVF).
5. Vaccination to protect the people from Cholera.

Hepatitis A

Definition - Hepatitis A is a systematic disorder that primarily affects the liver.

Causative organism

Heptatitis A virus (Entero virus)

Mode of transmission - Faeco oral route and direct contact

Incubation period- 14 -28 days

Clinical manifestations

1. Fever
2. Malaise
3. Severe anorexia, nausea and vomiting,
4. Pain in right hypochondriac region
5. Passing dark color urine and pale stool.

Complications

Acute liver failure

Control measures

a) Control of reservoir

Complete bed rest and disinfection of faeces and fomites by using 0.5 % sodium hypochlorite is recommended to prevent the spread of infection to others.

b) Control of transmission :

1. Hand washing before eating and after defaecation.
2. Sanitary disposal of excreta.

Unit_11_Communicable_disease_Gen_Nur.indd 250 09-03-2020 15:21:10